

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 142 877 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:
10.10.2001 Bulletin 2001/41

(21) Application number: 99961472.0

(22) Date of filing: 28.12.1999

(51) Int Cl.7: **C07D 215/48**, C07D 401/12,
C07D 405/12, C07H 15/203,
A61K 31/47, A61K 31/4709,
A61K 31/5355, A61K 31/706,
A61P 37/02, A61P 37/08,
A61P 29/00

(86) International application number:
PCT/JP99/07398

(87) International publication number:
WO 00/40562 (13.07.2000 Gazette 2000/28)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: 08.01.1999 JP 349899

(71) Applicant: **Japan Tobacco Inc.**
Tokyo 105-8422 (JP)


(72) Inventors:
• **INABA, Takashi**, Cent. Phar. Res. Inst. of Japan
Takatsuki-shi, Osaka 569-1125 (JP)

• **KAYA, Tetsudo**, Cent. Phar. Res. Inst. of Japan
Takatsuki-shi, Osaka 569-1125 (JP)
• **IWAMURA, Hiroyuki**, Cent. Phar. Res. of Japan
Takatsuki-shi, Osaka 569-1125 (JP)

(74) Representative: **VOSSIUS & PARTNER**
Siebertstrasse 4
81675 München (DE)

(54) 2-OXOQUINOLINE COMPOUNDS AND MEDICINAL USES THEREOF

(57) A 2-oxoquinoline compound or its pharmaceu-
tically acceptable salt of general formula [I]:



[I]

(wherein each symbol in the formula is as determined
in the description), and its pharmaceutical use.

The compound [I] of the present invention and its
pharmaceutically acceptable salts selectively act on
cannabinoid receptors, particularly on peripheral type
cannabinoid receptors, and have fewer side effects on
the central nervous system, having great immunosup-
pressive action, anti-inflammatory action or antiallergic
action. Therefore, these compounds are useful as can-
nabinoid receptors (particularly peripheral type cannab-
inoid receptors) modulator, immunosuppressants, anti-
inflammatory agents, and antiallergic agents.

EP 1 142 877 A1

DescriptionTechnical Field

5 **[0001]** The present invention relates to novel 2-oxoquinoline compounds and pharmaceutical uses thereof. In addition, the invention relates to novel uses of some types of 2-oxoquinoline compounds. More specifically, the invention relates to novel 2-oxoquinoline compounds that selectively act on cannabinoid receptors, particularly on peripheral cannabinoid receptors, and that have immunomodulating action, anti-inflammatory action, and antiallergic action with fewer side effects on the central nervous system and to pharmaceutical uses thereof.

Background Art

10 **[0002]** So far, a series of compounds named cannabinoids, which comprise C, H, and O, are known as constituents of cannabis. It has also been known that, among them, tetrahydrocannabinol (THC) is a major hallucinogen, and that a principal ingredient contained in the cannabis plant is Δ^9 -THC. It has been reported that Δ^9 -THC has actions including ataxia, increased sensitivity to stimulation, antiemetic action, analgesic action, hypothermic action, respiratory depression, catalepsy-inducing action, vasodilator action, immunosuppressive action, etc.

15 **[0003]** The sites of these actions of Δ^9 -THC are roughly categorized into the central nervous system (Devane et al., Mol Pharmacol. 1988, 34, 605-613; Hollister et al., Pharmacol. Rev., 1986, 38, 1-20; Renu et al., Prog. Drug Res., 1991, 36, 71-114) and the peripheral tissues (Nye et al., J. Pharmacol. Exp. Ther., 1985, 234, 784-791; Flynn et al., Mol Pharmacol. 1992, 41, 736-742). There are reports suggesting that some of the actions on the central nervous system are medically applicable.

20 **[0004]** On the other hand, it has been found that there exists the peripheral type receptor, for example, the receptor on macrophages (Munnro et al., Nature, 1993, 365, 61-65). Based on this finding, research and development are being carried out with the objective of designing peripheral type receptor agonists capable of modulating immune reaction and having anti-inflammatory action, antiallergic action, as well as the original immunomodulating action.


25 **[0005]** Further, agents selectively acting on peripheral type cannabinoid receptors can be safe agents that have no side effects on the central nervous system such as hypothermic action, catalepsy, and such, and therefore modulators selective for the peripheral type receptor are expected to be developed in particular.

30 **[0006]** Known cannabinoid receptor agonists include pyrazole derivatives (Unexamined Published Japanese Patent Application (JP-A) No. Hei 6-73014; EP Nos. 656354 and 658546), THC derivatives (JP-A Hei 3-209377), benzoxazine derivatives (US Patent No. 5,112,820), indole derivatives (US Patent No. 5,081,122), and aliphatic acid derivatives (WO 94/12466). However, there have previously been no findings on 2-oxoquinoline compounds by which the compounds of the present invention are characterized.


35 **[0007]** On the other hand, a variety of quinoline derivatives have been reported in terms of chemical structure.

40 **[0008]** For example, a reference of J. Pharm. Sci., 73, 11, 1652-1653 (1984) describes 6,7-dimethoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid benzylamide (compound A shown below), which is useful as a stimulant for the central nervous system. Another reference, Khim. Geterotsikl. Soedin., 8, 1101-1104 (1993), describes 4-hydroxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (pyridine-2-yl)amide (compound B shown below), which is useful as an anti-inflammatory agent. In Pharmaproject and others, it has been reported that 4-hydroxy-2-oxo-1-methyl-1,2-dihydroquinoline-3-carboxylic acid N-methyl-N-phenylamide (compound C shown below) that is also called "roquinimex" can be used as agents for a variety of diseases, e.g., anti-inflammatory agent, immunosuppressant, and antirheumatic agent.


45 **[0009]** Further, Examined Japanese Patent Publication No. Sho 47-14107 discloses a method for synthesizing 3-benzoylamino-6,7-dimethoxy-2-(1H)quinolone (compound D shown below) that can be used as an agent acting on the central nervous system.




Compound A



Compound B



Compound C (roquinimex)




Compound D


[0010] In addition, 2-oxo-1,2-dihydroquinoline-3-carboxylic acid cyclohexylamide has been disclosed in Synthesis, 11, 1362-1364 (1995). Quinoline compounds which are useful as immunomodulators have been disclosed in Published Japanese Translation of International Publication No. Hei 4-500373. Quinoline compounds which are useful as analgesic agents have been disclosed in French Patent Publication No. 2377400. Quinoline compounds which are useful as 5-HT₄ receptor agonists have been disclosed in republished patent publication WO 96/05166.

[0011] Further, one of 2-oxoquinoline compounds having 3,4-methylenedioxyphenyl group, which is one of specific features of the compounds of the present invention, is N-(3,4-methylenedioxyphenyl)-1,2-dihydro-4-hydroxy-1-methyl-2-oxoquinoline-3-carboxamide (compound E shown below), which is a compound capable of enhancing immunological activities and has been disclosed in JP-A Sho 57-171975 (EP No. 59698). Yet further, 1-methyl-2-oxo-3-{N-(1,3-benzodioxole-5-yl)-N-methylcarbamoyl}-4-hydroxy-6-methylthio-1,2-dihydroquinoline (compound F shown below), which has immunomodulating action, anti-inflammatory action, and analgesic action, has been disclosed in Published Japanese Translation of International Publication No. Hei 6-506925 (WO 92/18483).

[0012] However, these references have no description suggesting the existence of 2-oxoquinoline compounds of the present invention nor suggestion of pharmacological action based on the cannabinoid receptor-mediated mechanism.



Compound E




Compound F

[0013] On the other hand, description found in the publication of WO 97/29079 is that some compounds having quinoline structure, functioning as cannabinoid receptor agonist or antagonist, are useful as immunomodulators, anti-

inflammatory agents, antiallergic agents, or the like.


[0014] However, the quinoline compounds disclosed in the reference are merely quinolines that have been substituted with a hydroxy, and in the reference, there is no description suggesting the existence of the 2-oxoquinoline compounds of the present invention.

[0015] A reference of JP-A Hei 11-80124 (WO 99/02499) also describes immunomodulator, anti-inflammatory agent, and antiallergic agent comprising cannabinoid receptor agonist or antagonist as an active ingredient. The patent publication has also disclosed 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminophenyl) amide (compound G) and such, which is one of compounds having the 2-oxoquinoline structure that is one of specific features of the compounds of the present invention.



Compound G

[0016] In more detail, in the above-mentioned patent publication, for example, just merely the above-mentioned compound G and three other compounds are specifically disclosed, and the reference has no description of the inventive compounds that disclosed herein, which is represented by general formula [I], including compounds of which R^a is alkyl; compounds of which X is $-\text{COOR}^b$, $-\text{CONH}_2$, $-(\text{CH}_2)_p-\text{OC}(=\text{Y})-\text{NR}^d-(\text{Alk}^b)_s-\text{R}$, $-(\text{CH}_2)_q-\text{NR}^e-\text{C}(=\text{Z})-(\text{NR}^f)_w-(\text{Alk}^c)_v-\text{R}$, $-(\text{CH}_2)_p-\text{OH}$, or $-(\text{CH}_2)_q-\text{NR}^e\text{R}^e$ (where each symbol is as defined herein); compounds of which X is $-\text{CONR}^c-(\text{Alk}^a)_r-\text{R}$ and R^c is alkyl; or compounds of which R is represented by:



where each symbol is as defined herein.

[0017] It should be noted that, while the above-mentioned patent application was first laid open to public inspection on January 21, 1999 (WO 99/02499), the priority date of the present application is January 8, 1999, which is therefore earlier than the above publication date.

Disclosure of the Invention


[0018] An objective of the present invention is to provide novel compounds selectively acting on cannabinoid receptors, particularly on peripheral cannabinoid receptors and to provide pharmaceutical compositions thereof.

[0019] More specifically, the objective of the present invention is to provide novel compounds and pharmaceutical compositions thereof that are capable of selectively acting on cannabinoid receptors, particularly on the receptors of peripheral tissues, and that have therapeutic effects including immunomodulating action, anti-inflammatory action, antiallergic action, and such, but that hardly influence the central nervous system (specifically, side effects such as excitation, hallucination, ataxia, increased sensitivity to stimulation, hypothermic action, respiratory depression, catalepsy-inducing action, hypotension, etc.) and that are less toxic.

[0020] In order to achieve the above-mentioned objective, the present inventors strenuously studied and then found 2-oxoquinoline compounds exhibiting selective affinity for cannabinoid receptors, particularly for the peripheral type receptors, and thus being pharmaceutically useful for diseases associated with cannabinoid receptors, particularly diseases associated with peripheral type tissues (immune disease, various types of inflammation, allergic diseases, etc.). Thus, the inventors completed the present invention.

[0021] Specifically, the present invention comprises the following items (1)-(27):

(1) A cannabinoid receptor modulator comprising, as an active ingredient, a 2-oxoquinoline compound represented by the following general formula [I] or its pharmaceutically acceptable salt:



where W means -O-, -S(O)_t-, -CR³R⁴-, -NR⁵-, -NR⁵CO-, -CONR⁵-, -COO-, or -OCO- (where R³ and R⁴ may be identical or different and represent hydrogen atom or alkyl; R⁵ represents hydrogen atom or alkyl; and t indicates an integer, 0, 1 or 2);


R¹ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl; each group of R¹, except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, carboxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkylsulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R² represents hydrogen atom, alkyl, -OR⁶ (where R⁶ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl), -NR⁷R⁸ (where R⁷ and R⁸ may be identical or different and represent hydrogen atom, alkyl, alkenyl, alkynyl, acyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; or R⁷ and R⁸, together with the adjacent nitrogen atom, can form a heteroaryl); or -(CH₂)_u-S(O)_{u'}R⁹ (where R⁹ represents hydrogen atom, alkyl, alkenyl or alkynyl, each of u and u' independently represents an integer, 0, 1 or 2); each group of R², except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkyl sulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R^a represents hydrogen atom or alkyl;


X represents -COOR^b, -CONH₂, -CONR^c-(Alk^a)_r-R, -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R, -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R, -(CH₂)_p-OH or -(CH₂)_q-NR^eR^{e'}

(where each of R^b, R^c, R^d, and R^f independently represents hydrogen atom or alkyl; each of R^e and R^{e'} independently represents hydrogen atom or alkyl; or R^e and R^{e'}, together with the adjacent nitrogen atom, can form a heteroaryl; each of Alk^a, Alk^b and Alk^c independently represents alkylene or alkenylene; each of the alkylene and alkenylene may be substituted or unsubstituted with hydroxy, carboxy, alkoxycarbonyl, alkyl (the alkyl may be substituted or unsubstituted with hydroxy, alkoxy or alkylthio) or -CONR¹⁰R¹¹ (where R¹⁰ and R¹¹ may be identical or different and represent hydrogen atom or alkyl; or R¹⁰ and R¹¹, together with the adjacent nitrogen atom, can form a heteroaryl); R represents aryl, heteroaryl, cycloalkyl, benzene-condensed cycloalkyl or




where A and B independently represent oxygen atom, nitrogen atom or sulfur atom; k indicates an integer of 1-3; each of the aryl and heteroaryl may be substituted or unsubstituted with an alkyl substituted or unsubstituted with hydroxy or may be substituted or unsubstituted with a hydroxy, alkoxy, alkenyloxy, acyl, acyloxy, halogen atom, nitro, amino, sulfonamide, alkylamino, aralkyloxy, pyridyl, piperidino, carboxy, alkoxycarbonyl, acylamino, aminocarbonyl, cyano or glucuronic acid residue; the cycloalkyl may be substituted or unsubstituted with a hydroxy, alkoxy or =O; the benzene-condensed cycloalkyl may be substituted or unsubstituted with a hydroxy or alkoxy; each of r, s, v and w independently represents 0 or 1; each of Y and Z independently represents a nitrogen atom, oxygen atom or sulfur atom; and each of p and q independently represents an integer of 1-4).

(2) A cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of (1) or its pharmaceutically acceptable salt, wherein W represents -O-; R¹ is hydrogen atom or alkyl (the alkyl is as described above); R² represents -OR⁶ (R⁶ is as described above); and R is aryl, heteroaryl or



where aryl, heteroaryl, and each symbol in the formula are as defined above.

(3) A 2-oxoquinoline compound as represented by the following general formula [I'] or its pharmaceutically acceptable salt:



where W means -O-, -S(O)_t-, -CR³R⁴-, -NR⁵-, -NR⁵CO-, -CONR⁵-, -COO- or -OCO- (where R³ and R⁴ may be identical or different and represent hydrogen atom or alkyl; R⁵ represents hydrogen atom or alkyl; and t indicates an integer, 0, 1 or 2);


R¹ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; each group of R¹, except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, carboxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkylsulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R² represents hydrogen atom, alkyl, -OR⁶ (where R⁶ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl), -NR⁷R⁸ (where R⁷ and R⁸ may be identical or different and represent hydrogen atom, alkyl, alkenyl, alkynyl, acyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; or R⁷ and R⁸, together with the adjacent nitrogen atom, can form a heteroaryl); or - (CH₂)_u-S(O)_{u'}R⁹ (where R⁹ represents hydrogen atom, alkyl, alkenyl or alkynyl, each of u and u' independently represents an integer, 0, 1 or 2); each group of R², except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkyl sulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R^a represents hydrogen atom or alkyl;

X' represents -CONR^c-(Alk^a)_r-R, -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R or -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R, where each of R^c, R^d, R^e and R^f independently represents hydrogen atom or alkyl;

each of Alk^a, Alk^b and Alk^c independently represents alkylene or alkenylene; each of the alkylene and alkenylene may be substituted or unsubstituted with hydroxy, carboxy, alkoxycarbonyl, alkyl (the alkyl may be substituted or unsubstituted with a hydroxy, alkoxy or alkylthio); or -CONR¹⁰R¹¹ (where R¹⁰ and R¹¹ may be identical or different and represent hydrogen atom or alkyl; or R¹⁰ and R¹¹, together with the adjacent nitrogen atom, can form a heteroaryl); R represents aryl, heteroaryl, cycloalkyl, benzene-condensed cycloalkyl or




where A and B independently represent oxygen atom, nitrogen atom or sulfur atom; k indicates an integer of 1-3; each of the aryl and heteroaryl may be substituted or unsubstituted with an alkyl substituted or unsubstituted with hydroxy or may be substituted or unsubstituted with a hydroxy, alkoxy, alkenyloxy, acyl, acyloxy, halogen atom, nitro, amino, sulfonamide, alkylamino, aralkyloxy, pyridyl, piperidino, carboxy, alkoxycarbonyl, acylamino, amino-carbonyl, cyano or glucuronic acid residue; the cycloalkyl may be substituted or unsubstituted with a hydroxy, alkoxy or =O; the benzene-condensed cycloalkyl may be substituted or unsubstituted with a hydroxy or alkoxy; each of r, s, v and w independently represents 0 or 1; each of Y and Z independently represents a nitrogen atom,

oxygen atom or sulfur atom; and each of p and q independently represents an integer of 1-4, provided that:


- (a) when R² is a hydrogen atom, then WR¹ is substituted at position-j of 2-oxoquinoline; and
 (b) 1,2-dihydro-6,7-dimethoxy-2-oxo-N-(phenylmethyl)-3-quinolinecarboxamide and N-(1,2-dihydro-6,7-dimethoxy-2-oxo-3-quinolyl)benzamide are excluded.

- (4) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein R^a is alkyl.
 (5) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein X' is -CONR^c-(Alk^a)_r-R.
 (6) The 2-oxoquinoline compound of (5) or its pharmaceutically acceptable salt, wherein R^c is alkyl.
 (7) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein X' is -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R or -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R.
 (8) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein W is -O-; R¹ is hydrogen atom or alkyl having 1-3 carbons (the alkyl may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, carboxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkylsulfonyl); and R² is -OR⁶ {where R⁶ represents hydrogen atom or alkyl having 1-3 carbons (the alkyl may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkyl sulfonyl)}.
 (9) The 2-oxoquinoline compound of any one of (3)-(8) or its pharmaceutically acceptable salt, wherein R is aryl, heteroaryl or



where aryl, heteroaryl, and each symbol in the formula are as defined above.

- (10) The 2-oxoquinoline compound of any one of (3)-(8) or its pharmaceutically acceptable salt, wherein R is



where each symbol is as defined above.

- (11) The 2-oxoquinoline compound of any one of (3)-(10) or its pharmaceutically acceptable salt, wherein W is -O- and R² is -OR⁶ (where R⁶ is a hydrogen atom or alkyl).

- (12) The 2-oxoquinoline compound of any one of (3)-(11) or its pharmaceutically acceptable salt, wherein the substitution position of WR¹ is position-j of the benzene ring, and the substitution position of R² is position-i of the benzene ring.

- (13) The 2-oxoquinoline compound of any one of (3)-(6) and (8)-(12) or its pharmaceutically acceptable salt, wherein Alk^a is alkylene and r=1.

- (14) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound excludes 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide; 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminobenzyl)amide; 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-aminophenyl)ethyl]amide; and 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminophenyl)amide.

- (15) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound excludes compounds in which W is -O-; R¹ is alkyl; R² is -OR⁶ (where R⁶ is alkyl); R^a is hydrogen; and X' is -CONR^c-(Alk^a)_r-R (where R^c is a hydrogen atom; Alk^a is methylene, ethylene or trimethylene; r is 0 or 1; and R is aryl or heteroaryl).

- (16) The 2-oxoquinoline compound of (5) or its pharmaceutically acceptable salt, wherein R is aryl (the aryl may be substituted or unsubstituted with hydroxy, alkoxy, alkenyloxy, acyloxy, halogen atom, aralkyloxy, or glucuronic acid residue).

- (17) The 2-oxoquinoline compound of (3) or its pharmaceutically acceptable salt, wherein R is aryl except phenyl

or is heteroaryl except pyridyl.

(18) The 2-oxoquinoline compound of any one of (3)-(17) or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-1); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminobenzyl)amide (Example 3-2); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-aminophenyl)ethyl]amide (Example 3-3); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-4); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-5); 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-6); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide (Example 3-7); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-8); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-pyridyl methyl)amide (Example 3-9); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-piperidinoethyl)amide (Example 3-10); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide (Example 3-11); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3-pyridylmethyl)amide (Example 3-12); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridylmethyl)amide (Example 3-13); 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide (Example 3-14); 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-15); 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-16); 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide hydrochloride (Example 3-17); 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-18); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(2-fluorophenyl)ethyl]amide (Example 3-19); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(3-fluorophenyl)ethyl]amide (Example 3-20); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxy-3-methoxyphenyl)ethyl]amide (Example 3-21); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-chlorophenyl)ethyl]amide (Example 3-22); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide (Example 3-23); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-methylbenzyl)amide (Example 3-24); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-fluorobenzyl)amide (Example 3-25); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-26); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-27); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide (Example 3-28); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-29); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide (Example 3-30); 7,8-dimethoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-31); 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-32); 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-33); 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide (Example 3-34); 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-35); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-36); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide (Example 3-37); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide (Example 3-38); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-pyridylmethyl)amide (Example 3-39); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-fluorobenzyl)amide (Example 3-40); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide (Example 3-41); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-42); 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-43); 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide (Example 3-44); 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-45); 7,8-dipentyloxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 3-46); 8-hydroxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-47); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-dihydroxybenzyl)amide (Example 3-48); 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-hydroxy-3-methoxybenzyl)amide (Example 3-49); 1-O-{2-hydroxy-5-[(7-methoxy-2-oxo-8-pentyloxy-1,2-dihydro-3-quinolyl)carbonylamino methyl]phenyl}glucosidouronic acid and 1-O-{2-hydroxy-4-[(7-methoxy-2-oxo-8-pentyloxy-1,2-dihydro-3-quinolyl)carbonylamino methyl]phenyl}glucosidouronic acid (Example 3-50); 5-[7-methoxy-3-[(3,4-methylenedioxybenzyl)carbonyl]-2-oxo-1,2-dihydro-8-quinolyl]pentanoic acid (Exam-

ple 3-51); 5-[7-methoxy-3-((3-hydroxy-4-methoxybenzyl)carbamoyl)-2-oxo-1,2-dihydro-8-quinolyloxy]pentanoic acid (Example 3-52); 8-(5-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-53); 8-(5-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (4-hydroxy-3-methoxybenzyl)amide (Example 3-54); 8-(4-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-55); 7-methoxy-2-oxo-8-(4-oxopentyloxy)-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-56); 8-(3-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-57); 7-methoxy-2-oxo-8-(3-oxopentyloxy)-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-58); 8-(2-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (Example 3-59); 7,8-dihydroxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (Example 4-1); 8-butoxy-3-hydroxy methyl-7-methoxy-2-oxo-1,2-dihydroquinoline (Example 5-1); 8-ethoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (Example 5-2); N-(4-fluorophenyl)carbamic acid (8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)methyl ester (Example 6-1); N-pyridine-4-ylcarbamic acid (8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)methyl ester (Example 6-2); 3-dimethyl aminomethyl-8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline (Example 7-1); 8-butoxy-3-aminomethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (Example 7-2); 8-ethoxy-7-methoxy-3-morpholinomethyl-2-oxo-1,2-dihydroquinoline (Example 7-3); N-[(8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)methyl]-N'-(4-fluorophenyl)urea (Example 8-1) and N-[(8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)methyl]-(4-hydroxyphenyl)acetamide (Example 8-2).

(19) A 2-oxoquinoline compound or its pharmaceutically acceptable salt, the 2-oxoquinoline compound being selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-1); 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-2); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-3); and 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-4).

(20) A 2-oxoquinoline compound or its pharmaceutically acceptable salt, the 2-oxoquinoline compound being selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (Example 2-1); 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (Example 2-2); 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (Example 2-3); 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid (Example 2-4); 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (Example 2-5); 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (Example 2-6); and 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (Example 2-7).

(21) 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxamide (Example 3-60) or its pharmaceutically acceptable salt.

(22) A pharmaceutical composition comprising, as an active ingredient, the 2-oxoquinoline compound of any one of (3)-(21) or its pharmaceutically acceptable salt.

(23) A cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of any one of (3)-(21) or its pharmaceutically acceptable salt.

(24) A peripheral cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of any one of (3)-(21) or its pharmaceutically acceptable salt, the 2-oxoquinoline compound selectively acting on peripheral type cannabinoid receptors.

(25) The 2-oxoquinoline compound of any one of (3)-(21) or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is an immunomodulator, therapeutic agent for autoimmune diseases, antiallergic agent, or antiinflammatory agent.

(26) The 2-oxoquinoline compound of any of (3)-(21) or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is an antiinflammatory agent.

(27) An antiinflammatory agent comprising, as an active ingredient, the 2-oxoquinoline compound of (1) or (2) or its pharmaceutically acceptable salt.

[0022] Terminologies used herein are defined as follows:

[0023] The term "alkyl" means group having 1-10 carbon atoms, which may be linear or branched, and specifically includes methyl, ethyl, propyl, isopropyl, butyl, isobutyl, s-butyl, t-butyl, pentyl, isopentyl, neopentyl, t-pentyl, hexyl, isohexyl, neohexyl, heptyl, etc.

[0024] The group in R², R^{2'}, R⁵, and R⁶ preferably contains 1-7 carbon atoms; more preferred in R⁶ is a linear alkyl containing 1-6 carbon atoms and still more preferred in R⁶ is methyl. Preferably, the group in R³ and R⁴ has 1-4 carbon atoms. The group in R¹, R⁷, R⁸ and R⁹ is preferably a linear alkyl having 1-6 carbon atoms; more preferred is ethyl, propyl, butyl or pentyl; still more preferred is propyl, butyl or pentyl; and particularly preferred is pentyl. The group in R¹⁰, R¹¹, R¹² and R¹³ preferably has 1-4 carbon atoms; preferred in R^a, R^b, R^c, R^d, R^e, R^{e'} and R^f is 1-4 carbon atoms; and more preferred is methyl.

[0025] The term "alkenyl" means the linear or branched group having 2-10 carbon atoms, and is specifically exemplified by vinyl, allyl, crotyl, 2-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, heptenyl, etc.

[0026] The group in R⁶, R⁷, R⁸ and R⁹ preferably contains 2-7 carbon atoms. Preferably R¹ contains 4-7 carbon atoms.

[0027] The term "alkynyl" means a linear or branched group having 2-10 carbon atoms, and is specifically exemplified by ethynyl, propynyl, butynyl, 2-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, heptynyl, etc.

[0028] The group in R⁶, R⁷, R⁸ and R⁹ preferably contains 2-7 carbon atoms. Preferably R¹ contains 4-7 carbon atoms.

[0029] The "alkylene" in Alk^a, Alk^b and Alk^c means a linear or branched chain having 1-4 carbon atoms, which is specifically exemplified by methylene, ethylene, trimethylene, tetramethylene, etc., and more preferred is methylene or ethylene. Particularly preferred is methylene for Alk^a; and particularly preferred is methylene for each of Alk^b and Alk^c.

[0030] The "alkenylene" in Alk^a, Alk^b and Alk^c means a linear or branched chain having 2-4 carbon atoms, which is specifically exemplified by vinylene, propenylene, butenylene, etc.

[0031] The term "alkoxy" means the group of which alkyl portion corresponds to that having 1-4 carbon atoms among the above-defined alkyl groups, which is specifically exemplified by methoxy, ethoxy, propoxy, isopropoxy, butyloxy, t-butyloxy, etc.

[0032] The term "cycloalkyl" means a saturated monocyclic alkyl having 3-8 carbon atoms, which is specifically exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, etc.

[0033] Preferably the group in R¹, R⁶, R⁷ and R⁸ contains 3-6 carbon atoms; preferably R contains 3-7 carbon atoms and particularly preferred is cyclohexyl.

[0034] The "cycloalkylalkyl" in R¹, R⁶, R⁷ and R⁸ means the group in which the cycloalkyl portion is the above-defined cycloalkyl containing 3-6 carbon atoms and the alkyl portion is the above-defined alkyl containing 1-4 carbon atoms. Specifically, the group is exemplified by cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl, cyclopropylpropyl, cyclopropylbutyl, etc.

[0035] The "aryl" in R¹, R⁶, R⁷, R⁸ and R means an aromatic hydrocarbon having 6-16 carbon atoms and is specifically exemplified by phenyl, naphthyl, biphenyl, anthracenyl, indenyl, azulenyl, fluorenyl, phenanthrenyl, pyrenyl, etc.; preferred is phenyl or naphthyl; and particularly preferred is phenyl.

[0036] The "arylalkyl" in R¹, R⁶, R⁷, R⁸, R¹² and R¹³ means the group in which the aryl portion corresponds to the above-defined aryl and the alkyl portion is the above-defined alkyl having 1-4 carbon atoms. Specifically, the group is exemplified by benzyl, phenethyl, phenylpropyl, phenylbutyl, naphthylmethyl, biphenylmethyl, etc.; and preferred is benzyl.

[0037] The "heteroaryl" in R¹, R⁶, R⁷, R⁸ and R may be saturated or unsaturated with hydrogen atom and is specifically exemplified by pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, isooxazolyl, tetrazolyl, thiadiazolyl, oxadiazolyl, triazinyl, triazolyl, thienyl, pyrrolyl, pyrrolinyl, furyl, azepinyl, benzopyranlyl, benzothienyl, benzotriazolyl, indolyl, isoindolyl, quinolyl, isoquinolyl, cinnolyl, phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridyl, 1,7-naphthyridyl, 1,6-naphthyridyl, 1,5-naphthyridyl, pyrido[2,3-d]pyrimidyl, thieno[2,3-b]pyridyl, pyrrolidinyl, imidazolidinyl, piperidyl, piperazinyl, morpholyl, hydroazepinyl, hydroindolyl, hydroisoindolyl, hydroquinolyl, hydroisoquinolyl, etc.; preferred is pyridyl, thienyl, piperidyl, piperidino, imidazolyl, or morpholyl; more preferred is pyridyl, piperidyl, or morpholyl; and particularly preferred is pyridyl.

[0038] The expression, R^e and R^{e'} "together with the adjacent nitrogen atom, form a heteroaryl," means a heteroaryl having one or more nitrogen atoms among above heteroaryl groups; specifically, the group includes pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, morpholino, pyrazolyl, imidazolyl, tetrazolyl, triazolyl, pyrrolyl, pyrrolinyl, indolyl, hydroazepinyl, hydroindolyl, hydroisoindolyl, hydroquinolyl, hydroisoquinolyl, etc.; preferred is morpholino, piperidino or piperazinyl; and particularly preferred is morpholino.

[0039] The expression, R⁷ and R⁸, or R¹⁰ and R¹¹, "together with the adjacent nitrogen atom, form a heteroaryl" has the same meaning as the above-defined expression, R^e and R^{e'} "together with the adjacent nitrogen atom, form a heteroaryl."

[0040] The "heteroarylalkyl" in R¹, R⁶, R⁷ and R⁸ means the group in which the heteroaryl portion is the same as the above-defined one and the alkyl portion corresponds to the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by 2-thienylmethyl, 3-furylmethyl, 4-pyridylmethyl, 2-quinolylmethyl, 3-isoquinolylmethyl, etc.; and preferred is 4-pyridylmethyl.

[0041] The "benzene-condensed cycloalkyl" in R means the group of which cycloalkyl portion is the above-defined cycloalkyl; specifically, the group is exemplified by tetrahydronaphthalene, indan, etc.; and preferred is tetrahydronaphthalene.

[0042] The "acyl" in R⁷ and R⁸ means a group in which carbonyl has been substituted with the above-defined alkyl or the above-defined aryl; specifically, the group is exemplified by formyl, acetyl, propionyl, butyryl, valeryl, benzoyl, naphthoyl, etc.

[0043] Further, each of substituted or unsubstituted groups may be substituted or unsubstituted with one or more sub-

stituents, preferably with 1 or 2 substituents. Groups to be used as the substituents are described below.

[0044] The term "halogen atom" means fluorine, chlorine, bromine and iodine; and preferred are fluorine and chlorine.

[0045] The terms "alkyl", "alkoxy" and "acyl" indicate the same meanings as the above-defined "alkyl", "alkoxy" and "acyl," respectively.

[0046] The "alkoxycarbonyl" means the group of which alkyl portion is the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.; and preferred is ethoxycarbonyl.

[0047] The "alkylamino" means the group of which alkyl portion corresponds to the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by methylamino, ethylamino, propylamino, butylamino, dimethylamino, diethylamino, etc.

[0048] The "alkylthio" means the group of which alkyl portion is the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by methylthio, ethylthio, propylthio, butylthio, etc.

[0049] The "alkylsulfinyl" means the group of which alkyl portion corresponds to the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by methylsulfinyl, ethylsulfinyl, propylsulfinyl, butylsulfinyl, etc.

[0050] The "alkylsulfonyl" means the group of which alkyl portion is the above-defined alkyl having 1-4 carbon atoms; specifically, the group is exemplified by methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl, etc.

[0051] The "alkenyloxy" means the group of which alkenyl portion corresponds to the above-defined alkyl having 2-4 carbon atoms; specifically, the group is exemplified by ethenyloxy, propenyloxy, butenyloxy, etc.

[0052] The "acyloxy" means the group of which acyl portion is as defined above and is specifically exemplified by formyloxy, acetyloxy, propionyloxy, butyryloxy, isobutyryloxy, etc.; and preferred is acetyloxy.

[0053] The "acylthio" means the group of which acyl portion is as defined above and is specifically exemplified by formylthio, acetylthio, propionylthio, butyrylthio, isobutyrylthio, etc.; and preferred is acetylthio.

[0054] The "acylamino" means the group of which acyl portion is as defined above and is specifically exemplified by formylamino, acetylamino, propionylamino, butyrylamino, etc.; and preferred is acetylamino.

[0055] The "aralkyloxy" means the group of which aralkyl portion is the above-defined arylalkyl; and the group is specifically exemplified by benzyloxy, phenethyloxy, phenylpropyloxy, phenylbutyloxy, naphthylmethyloxy, biphenylmethyloxy, etc.

[0056] Preferred for R^a is hydrogen atom.

[0057] Substitution positions for the pair of WR¹ and R² are selected from the group consisting of position-g, position-h, position-i and position-j of 2-oxoquinoline but the respective substitution positions are different from each other. Preferred combination of the substitution positions is the combination of positions-h and -i, or that of positions-i and -j on the ring of 2-oxoquinoline; particularly preferred is the combination of positions-i and -j. It is preferable that R² is substituted at position-i.

[0058] Preferred for W is -O-, -S(O)_t- or -NR⁵-; more preferred is -O-. When W is -S(O)_t-, then t is preferred to be 0; when W is -NR⁵-, then it is preferable that R⁵ is a hydrogen atom.

[0059] Preferred for R¹ is a hydrogen atom or alkyl; more preferred is alkyl; and particularly preferred is unsubstituted alkyl.

[0060] Preferred substituents of alkyl include alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl and alkylsulfonyl; particularly preferred is hydroxy, carboxy or acyl. Alkyl, which has been substituted with any of hydroxy, carboxy, and acyl, is preferred to have 5 carbon atoms in total.

[0061] Specific examples of particularly preferred substituted alkyl include 2-hydroxypentyl, 3-hydroxypentyl, 4-hydroxypentyl, 5-hydroxypentyl, 2-oxopentyl, 3-oxopentyl, 4-oxopentyl, and 4-carboxybutyl.

[0062] Preferred for R² are the respective groups except hydrogen atom; specifically, such groups are alkyl, -OR⁶, -NR⁷R⁸ and -(CH₂)_u-S(O)_uR⁹ (where each symbol is as defined above); more preferred are -OR⁶, -NR⁷R⁸ and -(CH₂)_u-S(O)_uR⁹; and particularly preferred is -OR⁶.


[0063] When R² is -OR⁶, then preferred for R⁶ is hydrogen atom or alkyl; particularly preferred is alkyl. When R² is -NR⁷R⁸, then it is preferable that one of R⁷ and R⁸ is hydrogen atom and the other is alkyl. When R² is -(CH₂)_u-S(O)_uR⁹, preferred for u' and u is 0, and preferred for R⁹ is alkyl.

[0064] Preferred for X are -COOR^b, -CONR^c-(Alk^a)_r-R, -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R, -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R, -(CH₂)_p-OH and -(CH₂)_q-NR^eR^{e'} (where each symbol has the same meaning as defined above); particularly preferred is -COOR^b or -CONR^c-(Alk^a)_r-R; and more preferred for X and X' is -CONR^c-(Alk^a)_r-R; in this case, preferred for R^c is hydrogen atom.

[0065] When X and X' are -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R and -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R, then preferred for Y and Z is oxygen atom; preferred for p and q is 1; preferred for w is 1; preferred for s and v is 0; and preferred for R^e, R^d and R^f is hydrogen atom.

[0066] Preferred for R is aryl, heteroaryl or

Production method 1



In this formula, R¹² and R¹³ may be identical or different and represent hydrogen, alkyl, arylalkyl or cyano; other symbols

have the same meaning as defined above.

(Step 1)

[0076] Shown in this step is a method to obtain compound [3] by nitrating the ortho-position of formyl group on the benzene ring of compound [2].

[0077] It is possible to obtain a nitro compound by reacting compound [2] with fuming nitric acid in the presence of concentrated sulfuric acid in a solvent.

[0078] Such a solvent includes, for example, ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, etc.; and acid solvents such as acetic acid, acetic anhydride, and such; and preferred is acetic acid.

[0079] The reaction temperature is generally -50-200°C, and preferably -10-60°C. The reaction time is generally 15 minutes-48 hours, preferably 1-8 hours. The resulting nitro compound can further be reacted with an alkyl bromide such as bromopentane, etc. in the presence of a base in an adequate solvent to give compound [3].

[0080] The suitable base includes, for example, sodium carbonate, potassium carbonate, lithium carbonate, sodium bicarbonate, potassium bicarbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide, sodium hydride, n-butyllithium, s-butyllithium, t-butyllithium, and lithium diisopropylamide; and preferred is potassium carbonate.

[0081] The adequate solvent includes, for example, hydrocarbon solvents such as benzene, toluene, xylene, hexane, etc.; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; polar solvents such as dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, etc.; alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, and such; and preferred is dimethylformamide.

[0082] The reaction temperature is generally -10-200°C, preferably 0-60°C. The reaction time is generally 15 minutes-48 hours, preferably 1-8 hours.

(Step 2)

[0083] Compound [4] can be obtained by reducing the nitro group of compound [3] according to a commonly used method.

(Step 3)

[0084] Compounds [4] can be condensed with a malonic acid derivative [5] in the presence of an adequate acid or base to give compound [6]. The malonic acid derivative includes, for example, diethyl malonate, dimethyl malonate, dibenzyl malonate, ethyl cyanoacetate, methyl cyanoacetate, and such; and preferred is dimethyl malonate. The adequate acid includes, for example, benzoic acid, p-toluenesulfonic acid, acetic acid, methanesulfonic acid, hydrochloric acid, sulfuric acid, nitric acid, and such; benzoic acid is preferably used. The base includes, for example, sodium hydride, potassium t-butoxide, sodium ethoxide, sodium methoxide, ammonium acetate, sodium acetate, piperidine, pyridine, pyrrolidine, n-methylmorpholine, morpholine, triethylamine, and such; and preferred is piperidine.

[0085] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, xylene, hexane, heptane, etc.; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, etc.; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, etc.; polar solvents such as dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, etc.; and alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, etc.; and preferred is toluene.

[0086] The reaction temperature is generally 0-150°C, and preferably 120°C. The reaction time is generally 2-48 hours, and preferably 24 hours.

(Step 4)

[0087] Compound [6] can be hydrolyzed in the presence of an adequate base in a solvent to give compound [7].

[0088] The solvent includes alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, and such; water; or a mixed solvent thereof.

[0089] The suitable base includes, for example, sodium carbonate, potassium carbonate, lithium carbonate, sodium bicarbonate, potassium bicarbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide, sodium hydride, n-butyllithium, s-butyllithium, t-butyllithium, lithium diisopropylamide, and such; and preferred is lithium carbonate.

(Step 5)

[0090] Compound [7] that has been converted to an activated carboxylic acid derivative is allowed to react to compound [8] to yield compound [I] of interest.

5 [0091] The activated carboxylic acid derivative includes, for example, acid halide that can be obtained by treating carboxylic acid with thionyl chloride, phosphorus oxychloride, phosphorus pentachloride, oxalyl chloride, and such; active ester that can be obtained by condensing carboxylic acid with 1-hydroxybenzotriazole, N-hydroxysuccinimide or the like by using a condensing agent such as dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) hydrochloride, and such; mixed acid anhydride that can be obtained by reacting carboxylic acid to ethyl chlorocarbonate, pivaloyl chloride, isobutyl chlorocarbonate or the like, and such; preferably used is active ester that can be obtained from N-hydroxybenzotriazole by using EDC hydrochloride as a condensing agent.

10 [0092] If required, it is also possible to use a base in combination in the above reaction.

[0093] The base includes, for example, organic amines such as triethylamine, pyridine, and N-methylmorpholine; and preferred is triethylamine.

15 [0094] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, hexane, xylene, and such; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; polar solvents such as dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, and such; and preferred is dimethylformamide.

20 [0095] The reaction temperature is generally 0-100°C, and preferably 0-50°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

25

30


35

40


45

50




55

Production method 2

[0096] In this formula, R^x represents halogen atom; each of $R^{1'}$ and $R^{6'}$ independently represents alkyl; and R^b is as defined above.



In this formula, R^a represents alkyl; R^{1'}, R^{6'}, R^x, R^a and R^b are as defined above.



In this formula, R^{1''} represents alkyl; R^{1'}, R^{6'}, R^{7'}, R^a, R^c, Alk^a, R and r are as defined above.

35 (Step 1)

[0097] Compound [9] (3,4-dihydroxybenzaldehyde) is allowed to react by using alkylating agent [10] in the presence of a base, thereby selectively alkylating the hydroxyl group located at the 4-position of the phenol of compound [9]. This results in the production of compound [11].

40 [0098] The alkylating agent to be used includes alkyl iodide such as methyl iodide, etc.; alkyl bromide such as methyl bromide, ethyl bromide, propyl bromide, butyl bromide, pentyl bromide, and such; alkyl chloride such as pentyl chloride, and such; dialkyl sulfate such as dimethyl sulfate, and such; and preferably used is alkyl bromide.

45 [0099] The base includes, for example, sodium carbonate, potassium carbonate, lithium carbonate, cesium carbonate, calcium carbonate, sodium bicarbonate, potassium bicarbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide, sodium hydride, n-butyllithium, s-butyllithium, t-butyllithium, lithium diisopropylamide, and such; and preferably used is lithium carbonate.

50 [0100] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, xylene, hexane, and such; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; polar solvents such as dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, and such; alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, and such; and preferably used is dimethylformamide.

[0101] The reaction temperature is generally -20-100°C, and preferably 0-100°C. The reaction time is generally 15 minutes-48 hours, and preferably 1-6 hours.

55 (Step 2)

[0102] Compound [11] can be reacted with fuming nitric acid in the presence of concentrated sulfuric acid in a solvent,

thereby mono-nitrating the ortho-position of formyl group of compound [11] to obtain compound [12] (where compound [12] refers to compound [12a], compound [12b] and a mixture thereof).

[0103] The solvent includes, for example, ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, etc.; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; alcohol solvents such as methanol, ethanol, isopropyl alcohol, t-butanol, and such; acid solvents such as acetic acid, acetic anhydride, and such; preferably used is acetic acid.

[0104] The reaction temperature is generally -50-200°C, and preferably -10-60°C.

[0105] Alternatively, compound [12] can be obtained by nitrating compounds [11] by using lanthanum nitrate and sodium nitrate in the presence of an acid in a solvent in this step.

[0106] The acid includes, for example, benzoic acid, p-toluenesulfonic acid, acetic acid, methanesulfonic acid, hydrochloric acid, sulfuric acid, nitric acid, and such; and preferred is hydrochloric acid.

[0107] The solvent includes, for example, the above-mentioned solvents and preferred is tetrahydrofuran.

[0108] The reaction temperature is generally -50-200°C, and preferably 0-50°C.

(Step 3)

[0109] Compound [12] is allowed to react by using alkylating agent [13] in the presence of a base in a solvent, thereby alkylating the hydroxyl group located in the phenol of compound [12]. This results in the production of compound [14] (which means regioisomers and a mixture thereof corresponding to compound [12]).

[0110] The alkylating agent to be used includes the alkylating agents as described in Step 1 of Production method 2; and preferably used is alkyl bromide.

[0111] The base includes the bases as described in Step 1 of Production method 2; and preferably used is potassium carbonate.

[0112] The solvent includes the solvents as described in Step 1 of Production method 2; and preferably used is dimethylformamide.

[0113] The reaction temperature is generally -20-200°C, and preferably 0-100°C. The reaction time is generally 15 minutes-48 hours, and preferably 1-24 hr.

(Step 4)

[0114] Compound [14] is allowed to react to malonic acid in a solvent, thereby dehydration-condensing the formyl group moiety of compound [14]. This results in the production of compound [15] (which means regioisomers and a mixture thereof corresponding to compound [12]).

[0115] The solvent includes the solvents as described in Step 2 of Production method 2; and preferred is acetic acid.

[0116] The reaction temperature is generally -20-200°C, and preferably 0-100°C. The reaction time is generally 2-72 hours, but preferred is 3-24 hours.

(Step 5)

[0117] Compound [15] is esterified with alkylating agent [16] in the presence of a base in a solvent, thereby protecting the carboxyl group. This reaction results in the production of compound [17] (which means regioisomers and a mixture thereof corresponding to compound [15]).

[0118] The alkylating agent includes the alkylating agents as described in Step 1 of Production method 2; and preferred is methyl iodide.

[0119] The base includes the bases as described in Step 1 of Production method 2; and preferred is potassium carbonate.

[0120] The solvent includes the solvents as described in Step 1 of Production method 2; and preferred is dimethylformamide.

[0121] The reaction temperature is generally -20-200°C, and preferably 0-50°C. The reaction time is generally 15 minutes-48 hours, and preferably 1-24 hours.

(Step 6)

[0122] After the nitro group of compound [17] is reduced according to a commonly used method, the compound is condensed to construct the α -quinolone backbone. The reaction results in the production of compound [1-1a], compound [1-1a'], or a mixture thereof.

(Step 7)

[0123] Compound [I-1a], compound [I-1a'], or the mixture thereof, is alkylated by using alkylating agent [18] according to the same method as described in Step 3 of Production method 2. The reaction alkylates the NH portion of the quinolone moiety and this results in the production of the corresponding compound [I-1b], compound [I-1b'] or a mixture thereof.

(Step 8)

[0124] The esterified moiety of compound [I-1a], compound [I-1a'], compound [I-1b] or compound [I-1b'] is hydrolyzed according to a commonly used method to obtain the corresponding compound [I-2a], compound [I-2a'], compound [I-2b] and compound [I-2b'] (these four compounds are combined to be compound [I-2]).

(Step 9)

[0125] Compound [I-2], which has been converted to an activated carboxylic acid derivative, can be treated and amide-condensed with compound [8] in a solvent to give compound [I-3].

[0126] The activated carboxylic acid derivative includes, for example, acid halide that can be obtained by treating carboxylic acid with thionyl chloride, phosphorus oxychloride, phosphorus pentachloride, oxalyl chloride, and such; active ester that can be obtained by condensing carboxylic acid with 1-hydroxybenzotriazole, N-hydroxy succinimide or the like by using a condensing agent such as dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylamino propyl) carbodiimide (EDC) hydrochloride, and such; mixed acid anhydride that can be obtained by reacting carboxylic acid with ethyl chlorocarbonate, pivaloyl chloride, isobutyl chlorocarbonate or the like; and preferably used is active ester obtained from N-hydroxybenzotriazole by using EDC hydrochloride as a condensing agent.

[0127] If required, it is also possible to use a base in combination in the above reaction. The base includes, for example, organic amines such as triethylamine, pyridine, and N-methylmorpholine; and preferred is triethylamine.

[0128] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, hexane, xylene, and such; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; ester solvents such as ethyl acetate, methyl acetate, butyl acetate, and such; polar solvents such as dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, etc.; and preferred is dimethylformamide. The reaction temperature is generally 0-100°C, and preferably 0-50°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

(Step 10)

[0129] Compound [I-3] is treated with a Lewis acid in a solvent, thereby dealkylating the ether moiety. This reaction results in the production of compound [I-4].

[0130] The Lewis acid includes, for example, titanate tetrachloride, aluminum chloride, aluminum bromide, trimethylsilyl iodide, boron trichloride, boron tribromide, and such; and preferred is boron tribromide. Sulfur compound such as thiophenol, ethyl mercaptan, and such may be used in combination.


[0131] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, hexane, xylene, and such; ether solvents such as tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, etc.; and preferred is dichloromethane.

[0132] The reaction temperature is generally -100-100°C, and preferably -80-0°C. The reaction time is generally 15 minutes-24 hours, but preferred is 30 minutes-5 hours.

[0133] In this production method, it is also possible to obtain compound [I-3] by condensing compound [I-2a], [I-2a'] or a mixture thereof with compound [8] in the same manner as described in Step 9 of Production method 2 and then alkylating the NH portion of quinolone backbone in the same manner as described in Step 7 of Production method 2.

[0134] Alternatively, it is also possible to obtain compound [I-3], for example, by skipping Step 3, treating compound [12] by using the procedure of Step 4 and later procedures for the ring formation of 7-substituted 8-hydroxy-2-oxoquinoline, then carrying out the procedure of Step 9 for amide-condensation, followed by alkylation by the method of Step 3. Thus, different combinations of the steps can be used for the production.

Production method 3



[0138] If required, it is also possible to use a base in combination in the above reaction. The base includes, for example, organic amines such as triethylamine, pyridine, and N-methylmorpholine; and preferred is triethylamine. Used for the reduction of activated carboxylic acid are sodium borohydride, lithium borohydride, and such; and preferred is lithium borohydride.

[0139] The solvent includes, for example, hydrocarbon solvents such as benzene, toluene, hexane, xylene, etc.; ether solvents such as diethyl ether, 1,2-dimethoxyethane, tetrahydrofuran, diglyme, and such; halogen solvents such as dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, and such; and preferred is tetrahydrofuran.

[0140] In some cases in this reaction, base treatment should be carried out after reduction. The base being used in this reaction includes sodium hydroxide, potassium hydroxide, lithium hydroxide, potassium carbonate, sodium carbonate, and such; and preferred is sodium hydroxide. The reaction temperature is generally -20-100°C, and preferably -10-30°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

(Step 2)

[0141] Compound [I-5] is treated with isocyanate or thioisocyanate compound [20] in the presence of a base in a solvent to give compound [I-6].

[0142] The isocyanate and thioisocyanate include, for example, aralkyl isocyanate such as benzylisocyanate, and such; and arylisocyanate such as phenyl isocyanate, 4-fluorophenyl isocyanate, pyridine-4-yl isocyanate, and such; and preferred is arylisocyanate.

[0143] The base includes, for example, organic amines such as triethylamine, pyridine, N-methylmorpholine, and such; and preferred is triethylamine.

[0144] The solvent includes the solvents as described in Step 1 of Production method 2; and preferred is chloroform.

[0145] The reaction temperature is generally -20-100°C, and preferably -10-30°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

(Step 3)

[0146] Compound [I-5] is treated with acid chloride and then with amine compound [21] to give compound [I-7] in the presence of a base in a solvent.

[0147] The acid chloride includes methanesulfonyl chloride, p-toluenesulfonyl chloride, benzenesulfonyl chloride, and such; and preferred is methanesulfonyl chloride.

[0148] The base includes, for example, organic amines such as triethylamine, pyridine, N-methylmorpholine, and such; and preferred is triethylamine.

[0149] The solvent includes the solvents as described in Step 1 of Production method 2; and preferred is tetrahydrofuran.

[0150] The reaction temperature is generally -20-100°C, and preferably -10-30°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

(Step 4)

[0151] When at least R^e is a hydrogen atom in compound [I-7], compound [I-7] can be subjected to a coupling reaction with isocyanate or thioisocyanate compound [22] in a solvent to give the corresponding compound [I-8].

[0152] The isocyanate and thioisocyanate include those described in Step 1 of Production method 3; and preferably used is arylisocyanate.

[0153] If required, it is also possible to use a base in combination in the above reaction. The base includes, for example, organic amines such as triethylamine, pyridine, and N-methylmorpholine; and preferred is triethylamine.

[0154] The solvent includes the solvents as described in Step 1 of Production method 2; and preferred is dimethylformamide.

[0155] The reaction temperature is generally 0-100°C, and preferably 0-50°C. The reaction time is generally 15 minutes-24 hours, and preferably 1-12 hours.

(Step 5)

[0156] When at least R^e is a hydrogen atom in compound [I-7], compound [I-7] can be subjected to a coupling reaction with an activated carboxylic acid derivative in a solvent to obtain compound [I-8].

[0157] The carboxylic acid derivative to be used in the coupling reaction using an activated carboxylic acid derivative includes acid halide, active ester, mixed acid anhydride, etc, obtained from carboxylic acid [23] by the procedure as described in Step 9 of Production method 2; and preferred is an active ester obtained from N-hydroxybenzotriazole by

using EDC hydrochloride as a condensing agent.

[0158] The solvent includes the solvents as described in Step 1 of Production method 2; and preferred is dimethylformamide.

[0159] The reaction temperature is generally 0-100°C, and preferably 0-50°C. The reaction time is generally 15 minutes-24 hours, and preferably is 1-12 hours.

[0160] Compound [I] produced in the above-mentioned manner can be separated and purified, for example, by known methods such as concentration, concentration under reduced pressure, extraction with solvent, crystallization, re-crystallization, chromatography and the like.

[0161] Further, pharmaceutically acceptable salts of compound [I] and also various isomers of compound [I] can be produced according to previously known methods.

[0162] Compound [I] and pharmaceutically acceptable salts thereof exhibit pharmaceutical effects on diseases known to be associated with cannabinoid receptors, particularly diseases associated with peripheral cell tissues (immune disease, various types of inflammation, allergic diseases, etc.) in mammals.

[0163] In other words, compound [I] and pharmaceutically acceptable salts thereof selectively act on cannabinoid receptors, particularly on peripheral cannabinoid receptors, and thus have excellent immunomodulating action, anti-inflammatory action and antiallergic action, but exert fewer side effects on the central nervous system.

[0164] Thus, compound [I] and pharmaceutically acceptable salts thereof are useful as modulators for cannabinoid receptors (particularly for peripheral cannabinoid receptors), immunomodulators, therapeutic agents for autoimmune diseases, anti-inflammatory agents and antiallergic agents.

[0165] When the compound [I] or their pharmaceutically acceptable salts are used as a pharmaceutical composition, they may be formulated into tablets, pills, powders, granules, suppositories, injections, eye drops, liquid, capsules, troches, aerosols, elixirs, suspensions, emulsions, syrups, and so on by using a standard method, generally together with known pharmacologically acceptable carriers, excipients, diluents, extenders, disintegrators, stabilizers, preservatives, buffers, emulsifiers, aromatizers, colorants, edulcorants, viscosity increasing agents, flavors, solubilizers, or other additives such as water; plant oil; alcohols such as ethanol or benzyl alcohol, polyethylene glycol, glycerol triacetate, gelatin, lactose; or carbohydrates such as starch, magnesium stearate, talc, lanolin, vaseline, etc. The composition may be administered orally or parenterally.

[0166] The dose depends on the type and condition of disease, the type of compound to be administered and administration route, age, sex, and body weight of patient, etc. In general, compound [I] is daily given to an adult at a dose of 0.1-1000 mg, and preferably 1-300 mg, once or divided in the case of oral administration.

[0167] Further, the compounds of the present invention can be used as animal drugs.

Best Mode for Carrying out the Invention

[0168] The present invention is illustrated more specifically by referring to the following Examples. However, the present invention is not limited thereto.

Reference Example 1

4-Methoxy-2-nitro-3-pentyloxybenzaldehyde

[0169] Isovanillin (200 g), acetic acid (700 ml), and concentrated sulfuric acid (0.2 ml) were mixed and the resultant suspension was cooled to 0°C, to which a solution of fuming nitric acid (57.2 ml) in acetic acid (200 ml) was added dropwise over a period of 30 minutes. The mixture was stirred for 40 minutes, and then water (400 ml) was added thereto and the crystals were collected through filtration to give a mixture of 3-hydroxy-4-methoxy-2-nitrobenzaldehyde and 3-hydroxy-4-methoxy-6-nitrobenzaldehyde (56.4 g).

[0170] The resultant mixture was mixed with dimethylformamide (700 ml), and potassium carbonate (136.7 g) and bromopentane (127.7 ml) were successively added to this solution. After being stirred at 100°C for 4 hours, the reaction solution was filtered, and was separated with the addition of water (600 ml) and a mixture of hexane and ethyl acetate at a ratio of 1:1 (600 ml). The aqueous layer was extracted with a mixture of hexane and ethyl acetate at a ratio of 1:1 (600 ml), and the combined organic layer was dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The resultant crystals were collected through filtration. The filtrate was concentrated, and the resultant crystals were collected through filtration. This was repeated again, and the filtrate was concentrated to give 4-methoxy-2-nitro-3-pentyloxybenzaldehyde as a red oily substance (117 g). The filtered crystals were combined to give 4-methoxy-6-nitro-3-pentyloxybenzaldehyde as yellow crystals (90.1 g) (Table 1).

Reference Example 2

2-Amino-4-methoxy-3-pentyloxybenzaldehyde

5 **[0171]** 4-Methoxy-2-nitro-3-pentyloxybenzaldehyde (2.213 g) obtained in Reference Example 1 was dissolved in ethanol (22 ml), and tin chloride dihydrate (9.34 g) was added thereto. After being heat-refluxed for 4 hours, the reaction solution was cooled with ice, and a saturated aqueous solution of sodium hydrogencarbonate was added to the solution to render it alkaline. Then the solution was extracted with ethyl acetate (50 ml), and the organic layer was dried over
 10 anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The resultant residue was purified through silica-gel column chromatography (eluent: a mixture of n-hexane and ethyl acetate at a ratio of 5:1) to obtain 2-amino-4-methoxy-3-pentyloxybenzaldehyde (1.675 g) (Table 1).

Reference Example 3

3-Hydroxy-4-methoxybenzaldehyde

[0172] 3,4-Dihydroxybenzaldehyde (2.76 g, 20 mmol) was dissolved in DMF (15 ml), and methyl iodide (37.4 ml, 60 mmol) and anhydrous lithium carbonate (4.4 g, 60 mmol) were successively added to this solution. After being stirred at an external temperature of 90°C for 1.5 hours, the reaction mixture was cooled to room temperature and the inorganic
 20 salt was filtered off. A saturated aqueous solution of ammonium chloride (200 ml) was added to the filtrate to adjust the aqueous layer to a pH of from 7 to 8, and the solution was extracted with ethyl acetate (300 ml x 2). The combined organic layer was washed with a saturated aqueous solution of sodium chloride (300 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give the title compound (2.5 g, 82.2%) (Table 1).

Reference Example 4

3-Hydroxy-4-methoxy-2-nitrobenzaldehyde (4a)

3-Hydroxy-4-methoxy-6-nitrobenzaldehyde (4b)

[0173] 3-Hydroxy-4-methoxybenzaldehyde (15.2 g, 0.1 mol) obtained in the same manner as in Reference Example 3 was dissolved in THF (150 ml), to which sodium nitrate (8.5 g, 0.1 mol) and lanthanum nitrate hexahydrate (8.7 g, 0.2 mol) were added, and then a mixture of concentrated hydrochloric acid and water at a ratio of 1:1 (70 ml) was
 35 added dropwise over a period of 20 minutes. The mixture was stirred for 40 minutes at room temperature, and the separated organic layer was successively washed with water (50 ml), a saturated aqueous solution of sodium hydrogencarbonate (50 ml), and a saturated aqueous solution of sodium chloride (100 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give a mixture of the title compounds 4a and 4b as pale red crystals (4a:4b = 1:1, 13.3 g, yield 67.5%) (Table 2).

Reference Example 5

4-Methoxy-2-nitro-3-pentyloxybenzaldehyde (5a)

4-Methoxy-6-nitro-3-pentyloxybenzaldehyde (5b)

[0174] The mixture of 3-hydroxy-4-methoxy-2-nitrobenzaldehyde (4a) and 3-hydroxy-4-methoxy-6-nitrobenzaldehyde (4b) (4a:4b = 1:1, 12.3 g, 62.4 mmol) obtained in Reference Example 4 was dissolved in DMF (20 ml), and
 50 bromopentane (11.3 g, 74.9 mmol) and anhydrous potassium carbonate (12.9 g, 93.6 mmol) were successively added to this solution. After being stirred at an external temperature of 90°C for 1.5 hours, the reaction mixture was cooled to room temperature and the inorganic salt was filtered off. A saturated aqueous solution of ammonium chloride (200 ml) was added to the filtrate to adjust the aqueous layer to a pH of from 7 to 8, and the solution was extracted with ethyl acetate (300 ml x 2). The combined organic layer was washed with a saturated aqueous solution of sodium chloride (300 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated
 55 under reduced pressure to give a crude mixture of the title compounds 5a and 5b (5a:5b = 1:1, 17.12 g, quant.) (Tables 2 and 3). This crude product was used immediately in the subsequent reaction.

Reference Example 6

2-(4-Methoxy-2-nitro-3-pentyloxybenzylidene)malonic acid (6a)

2-(4-Methoxy-2-nitro-5-pentyloxybenzylidene)malonic acid (6b)

[0175] The crude product of 4-methoxy-2-nitro-3-pentyloxybenzaldehyde (5a) and 4-methoxy-6-nitro-3-pentyloxybenzaldehyde (5b) (5a:5b = 1:1, 17.12 g, 64.1 mmol) obtained in Reference Example 5 was dissolved in acetic acid (190 ml), to which malonic acid (20 g, 192 mmol) was added, and the mixture was stirred at an external temperature of 60°C for 16 hours. After toluene (100 ml x 2) was added to the reaction solution to remove acetic acid by azeotropic distillation, an aqueous solution of sodium hydroxide was added to the residue to adjust the aqueous solution to a pH of from 7 to 8, and the solution was washed with ethyl acetate (200 ml x 2). The aqueous layer was acidified (pH = 1 to 2) with concentrated hydrochloric acid, and was extracted with ethyl acetate (300 ml x 2). The combined organic layer was washed with a saturated aqueous solution of sodium chloride (300 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give a crude mixture of the title compounds 6a and 6b (6a:6b = 1:1, 23.3 g, overweight) (Table 3). This crude product was used immediately in the subsequent reaction.

Reference Example 7

Dimethyl 2-(4-methoxy-2-nitro-3-pentyloxybenzylidene)malonate (7a)

Dimethyl 2-(4-methoxy-2-nitro-5-pentyloxybenzylidene)malonate (7b)

[0176] The crude product of 2-(4-methoxy-2-nitro-3-pentyloxybenzylidene)malonic acid (6a) and 2-(4-methoxy-2-nitro-5-pentyloxybenzylidene)malonic acid (6b) (6a:6b = 1:1, 23.4 g, 64.1 mmol) obtained in Reference Example 6 was dissolved in DMF (160 ml), and methyl iodide (17.6 ml, 282 mmol) and anhydrous potassium carbonate (26.6 g, 192 mmol) were added to this solution. The mixture was stirred at room temperature for 1.5 hours, to which a saturated aqueous solution of ammonium chloride (800 ml) was added, and the solution was washed with ethyl acetate (300 ml x 2). The combined organic layer was successively washed with water (300 ml) and a saturated aqueous solution of sodium chloride (300 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give a crude mixture of the title compounds 7a and 7b (7a:7b = 1:1, 23.2 g, 94.8%) (Table 4). This crude product was used immediately in the subsequent reaction.

Reference Example 8

1-Methyl-4-bromo-1-t-butyl dimethylsilyl ether

Step 1

[0177] 2-Methyltetrahydrofuran (5.07 g, 41.7 mmol) was dissolved in trichloromethane (36 ml), and tetraethylammonium bromide (9.2 g, 43.8 mmol) was added thereto. Trifluoroborane etherate (5.56 ml, 43.8 mmol) was added dropwise to this solution at room temperature over a period of 10 minutes. After being stirred at room temperature for 16 hours, the reaction solution was cooled with ice, and a saturated aqueous solution of sodium hydrogencarbonate (40 ml) was added thereto to separate the organic layer. The aqueous layer was further extracted with trichloromethane (40 ml), and the combined organic layer was successively washed with water (40 ml) and a saturated aqueous solution of sodium chloride (40 ml), dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give 1-methyl-4-bromo-1-butanol (5.07 g, 72.8%) as a pale yellow oily substance. This product was used immediately in the subsequent reaction.

[0178] DMSO-d₆, 400 MHz: 4.4 (bs, 1H), 3.5-3.7 (m, 1H), 3.53 (t, 2H, J = 6.8 Hz), 1.8-1.9 (m, 2H), 1.3-1.5 (m, 2H), 1.04 (d, 3H, J = 6.2 Hz).

Step 2

[0179] 1-Methyl-4-bromo-1-butanol (5.07 g, 30.4 mmol) obtained in Step 1 was dissolved in dichloromethane (25 ml), and t-butyl dimethylsilyl chloride (9.64 g, 36.5 mmol) and imidazole (4.89 g, 45.6 mmol) were successively added thereto while being cooled with ice. After being stirred at room temperature for 7 hours, the reaction solution was cooled again with ice, and a saturated aqueous solution of sodium hydrogencarbonate (50 ml) was added thereto to separate

the organic layer. The organic layer was successively washed with water (50 ml) and a saturated aqueous solution of sodium chloride (50 ml), dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The residue was purified through silica-gel column chromatography (eluent: a mixture of hexane and ethyl acetate at a ratio of 100:1) to obtain the title compound (6.35 g., 74.3%).

[0180] DMSO-d₆, 400MHz: 3.8-3.9(m, 1H), 3.3-3.5(m, 2H), 1.8-2.0(m, 2H), 1.5-1.6(m, 2H), 1.13(d, 3H, J = 6.1 Hz), 0.87(s, 9H), 0.04(s, 6H).

Example 1-1

7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester

[0181] 2-Amino-4-methoxy-3-pentyloxybenzaldehyde (1.675 g) obtained in Reference Example 2 was dissolved in toluene (16 ml), to which dimethyl malonate (2.40 ml), piperidine (1.04 ml), and benzoic acid (80 ml) were added, and the mixture was stirred at an external temperature of 120°C for 27 hours. After the reaction solution was cooled to room temperature, a saturated aqueous solution of sodium hydrogencarbonate (1600 ml) was added thereto to separate the organic layer, and the aqueous layer was extracted with toluene (30 ml). The combined organic layer was washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The residue was purified through silica-gel chromatography (eluent: a mixture of chloroform and ethyl acetate at a ratio of 1:1) to obtain the title compound (251 mg) (Table 5).

Example 1-2 (Simultaneous synthesis with Example 1-1)

7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-1)

7-Methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (Example 1-2)

[0182] The crude product of dimethyl 2-[(4-methoxy-2-nitro-3-pentyl- oxyphenyl) methylene]propane-1,3-dioate (7a) and dimethyl 2-[(4-methoxy-2-nitro-5-pentyloxyphenyl)methylene]propane-1,3-dioate (7b) (7a:7b = 1:1, 23.1 g, 60.6 mmol) obtained in Reference Example 7 was dissolved in acetic acid (260 ml) and water (17 ml), and the solution was heated at an external temperature of 60°C. To this reaction solution, reduced iron (27.1 g, 48.5 mmol) was gradually added with attention to effervescence, and the mixture was further stirred at an external temperature of 90°C for 1.5 hours. After filtration, water (500 ml) was added to the filtrate, and the solution was extracted with ethyl acetate (300 ml x 2). The combined organic layer was successively washed with a 1% hydrochloric acid aqueous solution (500 ml) and a saturated aqueous solution of sodium chloride (300 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The residue was purified through column chromatography (eluent: a mixture of n-hexane and ethyl acetate at a ratio of 1:1) to obtain the title compounds of Example 1-1 (8.56 g, 44.2%) and Example 1-2 (4.83 g, 25.0%) as pale yellow crystals (Table 5). A crude product of 7a and 7b (7a:7b = 1:1, 23.2 g, 94.8%) was also obtained.

Example 1-3

1-Methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester

[0183] 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (1.02 g, 3.2 mmol) obtained in the same manner as in Example 1-1 was dissolved in DMF (10 ml), and methyl iodide (0.4 ml, 6.4 mmol) and anhydrous potassium carbonate (0.89 g, 6.4 mmol) were added to this solution. After the mixture was stirred at an external temperature of 60°C for 1.5 hours, a saturated aqueous solution of ammonium chloride (80 ml) was added thereto to adjust the solution to a pH of 8, and the solution was washed with ethyl acetate (50 ml x 2). The combined organic layer was successively washed with a saturated aqueous solution of sodium chloride (50 ml), dried over anhydrous magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The residue was purified through silica-gel chromatography (eluent: a mixture of chloroform and ethyl acetate at a ratio of 10:1) to obtain the title compound (532 mg, 49.9%) (Table 5).

[0184] A compound shown in Example 1-4 was obtained in the same manner as in Reference Examples and Example 1-3 above. Chemical structure and properties of the compound are shown in Table 6.

Example 2-1

7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid

[0185] 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester (240 mg) obtained in Example 1-1 was dissolved in methanol (7 ml) and water (3 ml), to which sodium hydroxide (120 ml) was added, and the solution was stirred at room temperature for 1.5 hours. While the reaction vessel was cooled with ice, the solution was acidified with the addition of concentrated hydrochloric acid, and extracted with ethyl acetate (20 ml). The organic layer was dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure to give the title compound (228 ml) (Table 6).

[0186] Compounds shown in Examples 2-2 to 2-7 were obtained in the same manner as in Reference Examples and Example 2-1 above. Chemical structures and properties of the compounds are shown in Tables 6 to 8.

Example 3-1

7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide

[0187] 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (30.0 mg) obtained in Example 2-1, 2-pyridine-4-ylethylamine (36.0 mg), and 1-hydroxybenzotriazole hydrate (17.3 mg) were dissolved in dimethylformamide (2 ml), and 1-ethyl-3-(3-dimethyl-aminopropyl)carbodiimide hydrochloride (24.5 mg) was successively added to this solution while being cooled with ice. After the reaction solution was stirred at room temperature for 5 hours, ethyl acetate (3 ml) and a saturated aqueous solution of sodium hydrogencarbonate (3 ml) were added to this solution. The organic layer was separated and dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The resultant residue was purified through column chromatography (eluent: a mixture of chloroform and methanol at a ratio of 25:1) to obtain the title compound (35 mg) as colorless crystals (Table 8).

[0188] Compounds shown in Examples 3-2 to 3-4 were obtained in the same manner as in Example 3-1 above. Chemical structures and properties of the compounds are shown in Table 9.

Example 3-5

7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide

[0189] 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (50 mg, 0.164 mmol) obtained in the same manner as in Example 2-1 was dissolved in dichloromethane (1 ml), and DMF (cat.) was added thereto, followed by thionyl chloride (0.018 ml, 0.24 mmol), and then the solution was stirred at room temperature for 1.0 hour. Toluene (4 ml x 2) was added to the reaction solution to remove the excess acid and thionyl chloride by evaporation, and an acid chloride was obtained as pale yellow crystals. The crystals were dissolved in dichloromethane (1 ml), and the solution was added dropwise to a solution of 3,4-methylenedioxybenzylamine (0.08 ml, 0.655 mmol) in dichloromethane (1 ml) prepared separately, and then the reaction solution was stirred at room temperature for 2 hours. The reaction solution was concentrated, and the residue was purified through column chromatography (eluent: a mixture of chloroform and ethyl acetate at a ratio of 10:1) to obtain the title compound (47 mg, 65.4%) (Table 10).

[0190] Compounds shown in Examples 3-6 to 3-54, and 3-60 were obtained in the same manner as in Reference Examples and Examples 3-1 and 3-5 above. Chemical structures and properties of the compounds are shown in Tables 10 to 26, and 28.

Example 3-55

8-(4-Hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide

[0191] In the usual manner, 8-hydroxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide (503 mg; Example 3-47) was alkylated with 1-methyl-4-bromo-1-*t*-butyldimethylsilyl ether obtained in Reference Example 8, and then *t*-butyldimethylsilyl group was removed to give the title compound (493 mg) (Table 26).

[0192] Compounds shown in Examples 3-56 to 3-59 were obtained in the same manner as in Example 3-55 above. Chemical structures and properties of the compounds are shown in Tables 27 and 28.

Example 4-1

7,8-Dihydroxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide

- 5 **[0193]** 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide (340 mg, 0.80 mmol) obtained in the same manner as in Example 3-5 was dissolved in dichloromethane (4 ml), and boron tribromide (1.0 M solution in dichloromethane; 2.4 ml, 2.4 mmol) was added dropwise thereto at an internal temperature of -60°C under a nitrogen stream. The solution was allowed to warm to room temperature, and was stirred for 0.5 hour. The reaction solution was poured into ice water (40 ml) to precipitate crystals. These crystals were collected through
10 filtration, washed with water (10 ml), and dried under reduced pressure to give the title compound (247.5 mg, 90.7%) (Table 29).

Example 5-1

- 15 8-Butoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline

[0194] 8-Butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (1.46 g, 5.00 mmol) obtained in the same manner as in Example 2-1 was dissolved in THF (20 ml), and triethylamine (1.53 ml, 11.0 mmol) and isopropyl chloro-
20 rocarbonate (1.35 g, 11.0 mmol) were successively added thereto while being cooled with ice. After the solution was stirred at the same temperature for 15 minutes, an aqueous solution of sodium borohydride (2.08 g, 55.0 mmol, 20 ml) was added thereto while being cooled with ice, and the solution was further stirred at the same temperature for 1 hour. A 50% sodium hydroxide aqueous solution (20 ml) was added to the solution, which was then stirred at room temper-
25 ature for 1 hour, and water (50 ml) and chloroform (50 ml) were added thereto to separate the organic layer. The aqueous layer was further extracted with chloroform (30 ml), and the combined organic layer was dried over magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The resultant residue was
25 purified through column chromatography (eluent: a mixture of hexane and ethyl acetate at a ratio of 1:3) to obtain the title compound (1.28 g, yield 92%) as colorless crystals (Table 29).

[0195] A compound shown in Example 5-2 was obtained in the same manner as in Example 5-1 above. Chemical
30 structure and properties of the compound are shown in Table 29.

Example 6-1

N-(4-Fluorophenyl)carbamic acid (8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl ester

- 35 **[0196]** 8-Butoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (100 mg, 0.361 mmol) obtained in the same manner as in Example 5-1 was dissolved in chloroform (1 ml), and triethylamine (10 µl) and 4-fluorophenyl isocyanate (45 µl, 0.396 mmol) were successively added thereto while being cooled with ice. The mixture was stirred at the same temperature for 5 hours, to which methanol (100 µl) was added, and concentrated under reduced pressure. The resultant residue was dissolved in THF, silica gel was slurried with the resultant solution, and the solution was
40 purified through column chromatography (eluent: a mixture of hexane and ethyl acetate at a ratio of 1:3) to obtain the title compound (128 mg, yield 86%) as colorless crystals (Table 30).

Example 6-2

- 45 N-Pyridine-4-ylcarbamic acid (8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl ester

[0197] 8-Ethoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (100 mg, 0.4 mmol) obtained in the same manner as in Example 5-1 was suspended in methylene chloride (1.5 ml), and pyridine (39 µl, 0.48 mmol) and 4-ni-
50 trophenylchloroformate (97 mg, 0.48 mmol) were added thereto while being cooled with ice. The mixture was stirred at the same temperature for 1 hour. After completion of the reaction, dimethylformamide (1.5 ml) was added, and then triethylamine (280 µl, 2.0 mmol) and 4-aminopyridine (188 mg, 2.0 mmol) were added to the reaction mixture while being cooled with ice, and the mixture was stirred at room temperature for 15 hours. After completion of the reaction, chloroform (15 ml) and a saturated aqueous solution of sodium bicarbonate (15 ml) were added to the mixture for
55 separation. The organic layer was dried over anhydrous sodium sulfate, filtered to remove the desiccating agent, and concentrated. The resultant residue was purified through preparative thin layer chromatography (developing eluent: a mixture of chloroform and methanol at a ratio of 90:10) to obtain the title compound (56 mg, yield 38%) as pale yellow crystals (Table 30).

Example 7-1

3-Dimethylaminomethyl-8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline

[0198] 8-Ethoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (200 mg, 0.80 mmol) obtained in the same manner as in Example 5-1 was dissolved in THF (5 ml), and triethylamine (134 μ l, 0.96 mmol) and methanesulfonyl chloride (68 μ l, 0.88 mmol) were successively added thereto while being cooled with ice. The mixture was stirred at the same temperature for 30 minutes, to which 40% aqueous solution of dimethylamine (7.0 ml) was added at a time, and was further stirred at room temperature for one hour. Water (20 ml) and chloroform (20 ml) were added to the reaction solution, and the organic layer was separated. The aqueous layer was further extracted with chloroform (10 ml), and the combined organic layer was concentrated. The resultant residue was dissolved in 1 N hydrochloric acid (5 ml), and the solution was washed three times with ethyl acetate (5 ml x 3). The aqueous layer was neutralized with a saturated aqueous solution of sodium bicarbonate, and was extracted two times with chloroform (5 ml x 2). The combined organic layer was dried over magnesium sulfate and concentrated under reduced pressure, and the resultant residue was washed with hexane to give the title compound (128 mg, yield 58%) as colorless crystals (Table 30).

Example 7-2

8-Butoxy-3-aminomethyl-7-methoxy-2-oxo-1,2-dihydroquinoline

[0199] 8-Butoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (431 mg, 1.55 mmol) obtained in the same manner as in Example 5-1 was dissolved in THF (10 ml), and triethylamine (260 μ l, 1.87 mmol) and methanesulfonyl chloride (132 μ l, 1.71 mmol) were successively added thereto while being cooled with ice. The mixture was stirred at the same temperature for 20 minutes, to which 28% aqueous ammonia (20 ml) was added at a time, and was further stirred at room temperature for 1 hour. Water (20 ml) and chloroform (20 ml) were added to the reaction solution, and the organic layer was separated. The aqueous layer was further extracted with chloroform (10 ml), and the combined organic layer was dried over magnesium sulfate, filtered to remove the desiccating agent, and concentrated under reduced pressure. The resultant residue was purified through column chromatography (eluent: a mixture of chloroform, methanol and aqueous ammonia at a ratio of 90:10:1) to obtain the title compound (146 mg, 34%) as colorless crystals (Table 31).

[0200] A compound shown in Example 7-3 was obtained in the same manner as in Example 7-1 or 7-2 above. Chemical structure and properties of the compound are shown in Table 31.

Example 8-1

N-[(8-Butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)-methyl]-N'-(4-fluorophenyl)urea

[0201] 8-Butoxy-3-aminomethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (56.6 mg, 0.205 mmol) obtained in Example 7-2 was dissolved in chloroform (1 ml), and 4-fluorophenyl isocyanate (25.6 μ l, 0.225 mmol) was successively added thereto while being cooled with ice. The mixture was stirred at the same temperature for 30 minutes, to which methanol (30 μ l) and triethylamine (30 μ l) were added, and was further stirred for 30 minutes. The reaction solution was concentrated under reduced pressure, and the resultant residue was washed with toluene to give the title compound (73 mg, yield 86%) as colorless crystals (Table 31).

Example 8-2

N-[(8-Butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-yl)-methyl]-(4-hydroxyphenyl)acetamide

[0202] 8-Butoxy-3-aminomethyl-7-methoxy-2-oxo-1,2-dihydroquinoline (77.7 mg, 0.281 mmol) obtained in Example 7-2, 4-hydroxyphenylacetic acid (47.0 mg, 0.309 mmol) and N-hydroxybenzotriazole (41.8 mg, 0.309 mmol) were dissolved in DMF (3 ml), and EDC (59.2 mg, 0.309 mmol) and triethylamine (30 μ l) were successively added thereto. The mixture was stirred at 50°C for 3 hours, and then a saturated aqueous solution of sodium bicarbonate (3 ml) and toluene (5 ml) were added thereto. The precipitated crystals were collected through filtration, and washed successively with water, 1 N hydrochloric acid, water, and ethyl acetate to give the title compound (88 mg, yield 76%) as colorless crystals (Table 32).

Table 1




Ref.Ex.No./structure/character/mp (°C)	1H NMR (δ) ppm	IR cm ⁻¹	MS
1 	CDCl ₃ , 300MHz 9.80(1H, s) 7.64(1H, d, J=8.6Hz) 7.09(1H, d, J=8.6Hz) 4.11(2H, t, J=6.6Hz) 3.99(3H, s) 1.60-1.80(2H, m) 1.28-1.47(4H, m) 0.92(3H, t, J=7.1Hz)		FAB+ 268[M+H] (80) 198 (100)
2 	CDCl ₃ , 300MHz 9.52(1H, s) 7.22(1H, d, J=9.0Hz) 6.90(1H, d, J=9.0Hz) 4.41(2H, t, J=6.9Hz) 3.97(3H, s) 2.0-2.3(2H, bs) 1.7-1.9(2H, m) 1.3-1.5(4H, m) 0.92(3H, t, J=7.2Hz)		FAB+ 238[M+H] (100)
3  Pale yellow crystal / 113-114	CDCl ₃ , 300MHz 9.84(s, 1H) 7.45(s, 1H) 7.43(d, 1H, J = 8.7 Hz) 6.98(d, 1H, J = 8.7 Hz) 6.40(bs, 1H) 4.00(s, 3H)		FAB+ 153[M+H] (100)

Table 2




Ref.Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
4a 	DMSO-d ₆ , 300MHz 10.91(bs, 1H) 9.57(s, 1H) 7.59(d, 1H, J=8.4Hz) 7.35(d, 1H, J=8.4Hz) 3.99(3H, s)		FAB- 232[M+H+] (20) 185(100)
4b 	DMSO-d ₆ , 300MHz 10.91(bs, 1H) 10.18(s, 1H) 7.83(s, 1H) 7.22(s, 1H) 3.96(3H, s)		FAB- 232[M+H+] (20) 185(100)
5a 	CDCl ₃ , 300MHz 9.79(s, 1H) 7.64(d, 1H, J = 8.6 Hz) 7.06(d, 1H, J = 8.6 Hz) 4.09(t, 2H, J = 6.7 Hz) 4.00(s, 3H) 1.7-1.8(m, 2H) 1.3-1.5(m, 4H) 0.92(t, 3H, J = 7.1 Hz)		

Table 3




Ref.Ex.No./structure/character/mp (°C)	1H NMR (δ) ppm	IR cm ⁻¹	MS
5b 	CDCl ₃ , 300MHz 10.43(s, 1H) 7.61(s, 1H) 7.39(s, 1H) 4.16(t, 2H, J = 6.7 Hz) 4.16(s, 3H) 1.8-2.0(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 6.9 Hz)		
6a 	CDCl ₃ , 300MHz 9.50(bs, 2H) 7.81(s, 1H) 7.38(d, 1H, J = 8.4 Hz) 7.00(d, 1H, J = 8.4 Hz) 4.05(t, 2H, J = 6.9 Hz) 3.93(s, 3H) 1.6-1.9(m, 2H) 1.3-1.5(m, 4H) 0.91(t, 3H, J = 6.9 Hz)		FAB+ 354[M+H+] (100)
6b 	CDCl ₃ , 300MHz 9.50(bs, 2H) 8.67(s, 1H) 7.76(s, 1H) 6.82(s, 1H) 4.10(t, 2H, J = 6.9 Hz) 3.97(s, 3H) 1.8-2.0(m, 2H) 1.3-1.5(m, 4H) 0.91(t, 3H, J = 6.9 Hz)		FAB+ 354[M+H+] (100)

Table 4



Ref.Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
7a  Pale blown crystal /	CDCl ₃ , 300 MHz 7.53(s, 1H) 7.19(d, 1H, J = 8.8 Hz) 6.97(d, 1H, J = 8.8 Hz) 4.04(t, 2H, J = 6.8 Hz) 3.93(s, 3H) 3.83(s, 3H) 3.79(s, 3H) 1.7-1.8(m, 2H) 1.3-1.5(m, 4H) 0.92(t, 3H, J = 7.0 Hz)		FAB+ 382[M-H+] (100)
7b  Pale blown crystal /	CDCl ₃ , 300 MHz 8.22(s, 1H) 7.76(s, 1H) 6.85(s, 1H) 4.11(t, 2H, J = 6.7 Hz) 3.97(s, 3H) 3.88(s, 3H) 3.66(s, 3H) 1.8-2.0(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 6.8 Hz)		FAB+ 382[M-H+] (100)

Table 5




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
1-1  Colorless crystal / 130-131	CDCl ₃ , 300MHz 9.04 (1H, bs) 8.50 (1H, s) 7.36 (1H, d, J=9.0Hz) 6.88 (1H, d, J=9.0Hz) 4.13 (2H, t, J=6.9Hz) 3.96 (3H, s) 3.74 (3H, s) 1.7-1.9 (2H, m) 1.3-1.5 (4H, m) 0.94 (3H, t, J=7.2Hz)	KBr 3423 2952 1742 1642 1505 1269	FAB+ 320(M+H+) (100) 288(70)
1-2  Colorless crystal / 194-195	CDCl ₃ , 300MHz 12.6 (s, 1H) 8.54 (s, 1H) 6.98 (s, 1H) 6.95 (s, 1H) 4.04 (t, 2H, J = 6.9 Hz) 4.04 (s, 3H) 3.70 (s, 3H) 1.8-2.0 (m, 2H) 1.3-1.6 (m, 4H) 0.95 (t, 3H, J = 6.9 Hz)	KBr 3444 2953 1730 1645 1571 1511 1484 1421 1269 1228 1900	FAB+ 320(M+H+) (100) 288(100)
1-3  Pale blown crystal / 102-103	CDCl ₃ , 300MHz 8.33(s, 1H) 7.36(d, 1H, J = 8.7 Hz) 6.92(d, 1H, J = 8.7 Hz) 3.98(s, 3H) 3.97(s, 3H) 3.94(s, 3H) 3.85(t, 2H, 6.8 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.93(t, 3H, J = 6.9 Hz)	KBr 2938 1703 1659 1613 1590 1500 1453 1274 1059 800	FAB+ 334[M-H+] (100)

Table 6




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
1-4  Colorless crystal / 124-125	DMSO-d ₆ , 300MHz 8.42(s, 1H) 7.44(s, 1H) 6.97(s, 1H) 3.98(t, 2H, J = 6.6 Hz) 3.97(s, 3H) 3.78(s, 3H) 3.65(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.90(t, 3H, J = 6.8 Hz)	KBr 2948 1736 1260 1080 795	FAB+ 334(M+H+) (100) 302(100)
2-1  Pale yellow crystal / 178-179	CDCl ₃ , 300MHz 14.00(1H, s) 9.42(1H, bs) 8.87(1H, s) 7.51(1H, d, J=9.0Hz) 7.03(1H, d, J=9.0Hz) 4.18(2H, t, J=6.9Hz) 4.01(3H, s) 1.7-1.8(2H, m) 1.3-1.5(4H, m) 0.95(3H, t, J=6.9Hz)	KBr 3423 2952 1742 1642 1505 1269	FAB+ 306[M+H+] (100)
2-2  Colorless crystal / 182-184	CDCl ₃ , 300MHz 14.98 (br s, 1H) 9.51 (br s, 1H) 8.87 (s, 1H) 7.51 (d, J=9.0 Hz, 1H) 7.04 (d, J=9.0 Hz, 1H) 4.20 (t, J=6.9 Hz, 2H) 4.02 (s, 3H) 1.81 (m, 2H) 1.51 (m, 2H) 1.00 (t, J=6.9 Hz, 3H)	KBr 1723 1630 1506 1283 1258 1099	

Table 7




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
2-3  Colorless crystal / 242-244	CDCl ₃ , 300MHz 14.00 (br s, 1H) 9.71 (br s, 1H) 8.88 (s, 1H) 7.52 (d, J=9.0 Hz, 1H) 7.04 (d, J=9.0 Hz, 1H) 4.29 (q, J=6.9 Hz, 2H) 4.02 (s, 3H) 1.44 (t, J=6.9 Hz, 3H)	KBr 1736 1634 1504 1474 1282 1259 1095	
2-4  Colorless crystal / 178-181	DMSO-d ₆ , 300MHz 12.37 (brs, 1H) 8.90 (s, 1H) 7.81 (d, 1H, J=9.2Hz) 7.27 (d, 1H, J=9.2Hz) 3.97 (m, 5H) 1.80 (q, 2H, J=7.3Hz) 0.95 (t, 3H, J=7.6Hz)	KBr 3188 1735 1630 1507 1286	
2-5  Colorless crystal / 290<dec.	DMSO-d ₆ , 300MHz 14.96 (s, 1H) 13.05 (s, 1H) 8.81 (s, 1H) 7.55 (s, 1H) 7.01 (s, 1H) 4.01 (t, 2H, J = 6.6 Hz) 3.90 (s, 3H) 1.7-1.9 (m, 2H) 1.3-1.5 (m, 4H) 0.91 (t, 3H, J = 6.8 Hz)	KBr 3422 1685 1211	FAB+ 305 (M+H+)(100)

Table 8




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
2-6  Colorless crystal / 188-189	DMSO-d ₆ ,300MHz 14.58(s, 1H) 8.76(s, 1H) 7.52(d, 1H, J = 8.8 Hz) 7.07(d, 1H, J = 8.8 Hz) 4.02(s, 3H) 3.93(s, 3H) 3.91(t, 2H, J = 6.9 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 2956 1735 1622 1560 1509 1458 1379 1280 1067	FAB+ 320(M+H+) (100) 302(100)
2-7  Colorless crystal / 196-197	CDCl ₃ ,300MHz 14.70(s, 1H) 8.78(s, 1H) 7.11(s, 1H) 6.83(s, 1H) 4.08(t, 2H, J = 6.8 Hz) 4.06(s, 3H) 3.86(s, 3H) 1.8-2.0(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 3448 1718 1654 1560 1508 1271	FAB+ 320(M+H+) (50) 302(100)
3-1  Colorless crystal / 98.0~99.0°C	CDCl ₃ ,300MHz 9.74(1H,br) 9.13(1H,bs) 8.49(1H,s) 8.53(2H,d,J=6.0Hz) 7.46(1H,d,J=8.9Hz) 7.22(2H,d,J=6.0Hz) 6.94(1H,d,J=8.9Hz) 4.14(2H,t,J=6.9Hz) 3.98(3H,s) 3.76(2H,q,J=6.7Hz) 2.97(2H,t,J=7.2Hz) 1.74-1.88(2H,m) 1.35-1.53(4H,m) 0.95(3H,t,J=7.1Hz)	KBr 3257 2938 1672 1622 1530 1261 1112 805	FAB+ 410[M+H+] (60) 288 (60)

Table 9




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-2  Colorless amorphous /	CDCl ₃ , 300MHz 9.87(1H, br) 9.10(1H, bs) 8.89(1H, s) 7.45(1H, d, J=8.8Hz) 7.18(2H, d, J=8.3Hz) 6.93(1H, d, J=8.8Hz) 6.66(2H, d, J=8.3Hz) 4.56(2H, d, J=6.0Hz) 4.13(2H, t, J=6.9Hz) 3.97(3H, s) 1.71-1.87(2H, m) 1.30-1.50(4H, m) 0.94(3H, t, J=7.1Hz)	KBr 3232 2954 1668 1622 1520 1260 1109 801	FAB+ 410[M+H+] (40) 288 (20)
3-3  Colorless crystal / 125.0~126.0°C	CDCl ₃ , 300MHz 9.63(1H, br) 9.10(1H, bs) 8.85(1H, s) 7.44(1H, d, J=8.7Hz) 7.07(2H, d, J=8.4Hz) 6.93(1H, d, J=8.7Hz) 6.66(1H, d, J=8.4Hz) 4.13(2H, t, J=6.8Hz) 3.97(3H, s) 3.67(2H, q, J=6.8Hz) 2.83(2H, t, J=7.2Hz) 1.75-1.88(2H, m) 1.30-1.50(4H, m) 0.95(3H, t, J=7.2Hz)	KBr 3248 2928 1672 1625 1539 1260 1112 802	FAB+ 424[M+H+] (40)
3-4  Colorless crystal / 267.0~268.0°C	DMSO-d ₆ , 300MHz 12.1(1H, s) 11.7(1H, s) 8.88(1H, s) 7.77(2H, d, J=9.0Hz) 7.75(1H, d, J=9.0Hz) 7.26(2H, d, J=9.0Hz) 7.18(1H, d, J=9.0Hz) 4.00(2H, t, J=6.9Hz) 1.70-1.83(2H, m) 1.30-1.50(4H, m) 0.89(3H, t, J=7.2Hz)	KBr 2954 1672 1622 1552 1498 1284 1263	FAB+ 296 (30) 288 (100) 218 (70)

Table 10




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-5  Colorless crystal / 126-127	CDCl ₃ ,300MHz 13.99 (br t, 1H) 9.18(s, 1H) 8.90(s, 1H) 7.47(d, 1H, J = 8.8 Hz) 6.95(d, 1H, J = 8.8 Hz) 6.8-6.9(m, 2H) 6.77(d, 1H, J = 7.9 Hz) 5.95(s, 2H) 4.59(d, 2H, J = 5.8 Hz) 4.14(t, 2H, J = 8.7 Hz) 3.99(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 2956 1684 1619 1535 1263	FAB+ 439[M+H+] (100)
3-6  Colorless crystal / 201-211	CDCl ₃ ,300MHz 9.80 (br t, J=5.9 Hz, 1H) 9.21 (br s, 1H) 8.81 (s, 1H) 8.58 (d, J=5.9 Hz, 2H) 7.53 (d, J=6.2 Hz, 2H) 7.47 (d, J=8.8 Hz, 1H) 6.96 (d, J=8.8 Hz, 1H) 4.25 (q, J=7.1 Hz, 2H) 3.99 (s, 3H) 3.83 (q, J=6.6 Hz, 2H) 3.12 (t, J=7.0 Hz, 2H) 1.43 (t, J=7.1 Hz, 3H)	KBr 1672 1626 1538 1500 1375 1286 1260 1223 1115	(ESI+) 368 264 246 (ESI-) 366 323
3-7  Colorless crystal / 106-107	9.71(br, 1H) 9.18(s, 1H) 8.86(s, 1H) 7.44(d, 1H, J = 8.7 Hz) 7.12(d, 2H, J = 8.4 Hz) 6.93(d, 1H, J = 8.4 Hz) 6.80(d, 2H, J = 8.4 Hz) 6.23(s, 1H) 4.12(t, 2H, J = 6.6 Hz) 3.97(s, 3H) 3.70(q, 2H, J = 7.2 Hz) 2.87(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.2 Hz)	KBr 3246 2932 1673 1625 1537 1515 1500 1262 1110	FAB+ 425(M+H+) (100) 288(75)

Table 11




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-8  Colorless crystal / 105-106	9.67(bt, 1H) 9.11(s, 1H) 8.85(s, 1H) 7.45(d, 1H, J = 8.7 Hz) 7.2-7.3(m, 2H) 7.00(t, 2H, J = 8.7 Hz) 6.93(d, 1H, J = 8.7 Hz) 4.13(t, 2H, J = 6.9 Hz) 3.97(s, 3H) 3.97(q, 2H, J = 7.2 Hz) 2.92(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.6(m, 4H) 0.95(t, 3H, J = 6.9 Hz)	KBr 3286 2962 1661 1614 1533 1497 1262	FAB+ 427(M+H+) (100) 288(40)
3-9  Colorless crystal / 164-163	CDCl ₃ , 300MHz 10.16(bt, 1H) 9.26(s, 1H) 8.90(s, 1H) 8.55(d, 2H, J = 6.8 Hz) 7.47(d, 1H, J = 8.8 Hz) 7.28(d, 2H, J = 6.8) 6.87(d, 1H, J = 8.8 Hz) 4.70(t, 2H, J = 6.0 Hz) 4.15(t, 2H, J = 6.9 Hz) 3.99(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 6.9 Hz)	KBr 2956 1684 1619 1535 1263	FAB+ 396(M+H+) (100) 288(40)
3-10  Yellow oil /	CDCl ₃ , 300MHz 9.76(bt, 1H) 9.12(s, 1H) 8.85(s, 1H) 7.44(d, 1H, J = 9.0 Hz) 6.93(d, 1H, J = 9.0 Hz) 4.13(t, 2H, J = 6.9 Hz) 3.97(s, 3H) 3.60(q, 2H, J = 6.6 Hz) 2.59(t, 2H, J = 6.6 Hz) 2.4-2.5(m, 4H) 1.7-1.9(m, 2H) 1.5-1.7(m, 4H) 1.3-1.5(m, 6H) 0.95(t, 3H, J = 6.9 Hz)	Near 2932 1667 1537 1504	FAB+ 416(M+H+) (100) 331(50) 288(40)

Table 12




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-11  Colorless crystal / 94-96	CDCl ₃ ,300MHz 9.81(bt, 1H) 9.13(s, 1H) 8.86(s, 1H) 7.45(d, 1H, J = 8.9 Hz) 6.94(d, 1H, J = 8.9 Hz) 4.13(t, 2H, J = 6.9 Hz) 3.98(s, 3H) 3.7-3.8(m, 4H) 3.61(q, 2H, J = 6.4 Hz) 2.62(t, 2H, J = 6.4 Hz) 2.5-2.6(m, 4H) 1.7-1.9(m, 2H) 1.4-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 3438 2953 1718 1639 1507 1486 1287	FAB+ 418(M+H+) (100)
3-12  Colorless crystal / 143-144	CDCl ₃ ,300MHz 10.07(bt, 1H) 9.16(s, 1H) 8.89(s, 1H) 8.64(s, 1H) 8.52(d, 1H, J = 1.7 Hz) 7.7(d, 1H, J = 7.9 Hz) 7.47(t, 1H, J = 8.8 Hz) 7.26(dd, 1H, J = 7.9, 1.7 Hz) 6.95(d, 1H, J = 8.8 Hz) 4.70(d, 2H, J = 6.0 Hz) 4.14(t, 2H, J = 6.9 Hz) 3.98(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.1 Hz)	KBr 2956 1684 1618 1537 1264	FAB+ 396(M+H+) (100) 288(40)
3-13  Colorless crystal / 141-142	CDCl ₃ ,300MHz 10.35(bt, 1H) 9.17(s, 1H) 8.90(s, 1H) 8.60(d, 1H, J = 4.7 Hz) 7.65(dt, 1H, J = 7.7, 1.7 Hz) 7.46(d, 1H, J = 8.9 Hz) 7.61(t, 1H, J = 7.7 Hz) 7.16(dt, 1H, J = 4.7, 1.7 Hz) 6.94(d, 1H, J = 8.9 Hz) 4.84(t, 2H, J = 5.6 Hz) 4.14(t, 2H, J = 6.9 Hz) 3.98(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 6.9 Hz)	KBr 2931 1668 1622 1526 1262	FAB+ 396(M+H+) (100) 288(40)

Table 13




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-14  Colorless crystal / 91-93	CDCl ₃ ,300MHz 9.97 (br t, J=5.9 Hz, 1H) 9.12 (br s, 1H) 8.86 (s, 1H) 7.45 (d, J=8.8 Hz, 1H) 7.35-7.23 (m, 5H) 6.93 (d, J=8.8 Hz, 1H) 4.14 (t, J=7.0 Hz, 2H) 3.97 (s, 3H) 3.74 (q, J=6.8 Hz, 2H) 2.95 (t, J=7.1 Hz, 2H) 1.81 (quintet, J=7.1 Hz, 2H) 1.51 (m, 2H) 1.00 (t, J=7.3 Hz, 3H) br t, J=5.9 Hz, 1H	KBr 1660 1613 1535 1375 1260 1133	(fab+, NBA) 395 274 218 (fab-, NBA) 393 305 153
3-15  Colorless crystal / 125-127	CDCl ₃ ,300MHz 9.67 (br t, J=5.9 Hz, 1H) 9.12 (br s, 1H) 8.85 (s, 1H) 7.45 (d, J=8.8 Hz, 1H) 7.24 (m, 2H) 7.00 (m, 2H) 6.93 (d, J=8.8 Hz, 1H) 4.14 (t, J=7.0 Hz, 2H) 3.97 (s, 3H) 3.71 (q, J=6.8 Hz, 2H) 2.92 (t, J=7.1 Hz, 2H) 1.80 (quintet, J=7.1 Hz, 2H) 1.51 (m, 2H) 1.00 (t, J=7.3 Hz, 3H)	2964 1661 1613 1532 1376 1259 1111	(fab+, NBA) 413 274 218 (fab-, NBA) 411 305 153
3-16  Colorless crystal / 135-136	CDCl ₃ ,300MHz 9.73 (br t, J=5.9 Hz, 1H) 9.14 (br s, 1H) 8.85 (s, 1H) 8.53 (d, J=6.2 Hz, 2H) 7.45 (d, J=8.8 Hz, 1H) 7.21 (d, J=5.9 Hz, 2H) 6.94 (d, J=8.8 Hz, 1H) 4.15 (t, J=6.9 Hz, 2H) 3.98 (s, 3H) 3.76 (q, J=6.7 Hz, 2H) 2.96 (t, J=7.3 Hz, 2H) 1.80 (quintet, J=7.5 Hz, 2H) 1.51 (m, 2H) 1.00 (t, J=7.3 Hz, 3H)	KBr 1676 1626 1537 1499 1370 1284 1260 1113	(fab+, NBA) 396 274 218 (fab-, NBA) 394 305 153

Table 14




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-17  Colorless crystal / 218 (decomp)	DMSO-d ₆ , 300MHz (br s, 1H) 9.79 (t, J=5.9 Hz, 1H) 8.84 (d, J=6.6 Hz, 2H, J=6.6 Hz) 8.72 (s, 1H) 8.00 (d, J=6.6 Hz, 2H) 7.68 (d, J=8.8 Hz, 1H) 7.13 (d, J=8.8 Hz, 1H) 3.99 (t, J=6.9 Hz, 2H) 3.93 (s, 3H) 3.75 (q, J=6.5 Hz, 2H) 3.19 (t, J=6.6 Hz, 2H) 1.76 (quintet, J=7.3 Hz, 2H) 1.41 (m, 2H) 0.93 (t, J=7.3 Hz, 3H)	KBr 1676 1626 1537 1499 1370 1284 1260 1113	(ESI+) 396 292 274 (ESI-) 394 338 322
3-18  Colorless crystal / 172-173	CDCl ₃ , 300MHz 9.67 (br t, J=5.9 Hz, 1H) 9.17 (br s, 1H) 8.86 (s, 1H) 7.46 (d, J=8.8 Hz, 1H) 7.24 (m, 2H) 7.00 (m, 2H) 6.94 (d, J=8.8 Hz, 1H) 4.23 (q, J=7.1 Hz, 2H) 3.98 (s, 3H) 3.70 (q, J=7.1 Hz, 2H) 2.92 (t, J=7.1 Hz, 2H) 1.42 (t, J=7.0 Hz, 3H)	KBr 1666 1626 1509 1262 1218 1114	(ESI+) 385 264 246 (ESI-) 383 355 341
3-19  Colorless crystal / 115-116	CDCl ₃ , 300MHz 9.65 (br, 1H) 9.12 (s, 1H) 8.82 (s, 1H) 7.40 (d, 1H, J = 8.8 Hz) 6.9-7.3 (m, 4H) 6.90 (d, 1H, J = 8.8 Hz) 4.09 (t, 2H, J = 6.6 Hz) 3.93 (s, 3H) 3.67 (q, 2H, J = 7.0 Hz) 2.96 (t, 2H, J = 7.0 Hz) 1.7-1.9 (m, 2H) 1.3-1.5 (m, 4H) 0.91 (t, 3H, J = 7.0 Hz)	KBr 3235 2951 1663 1611 1530 1483 1286	FAB+ 427 (M+H+) (100)

Table 15




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-20  Colorless crystal / 115-116	9.69(bt, 1H) 9.16(s, 1H) 8.86(s, 1H) 7.45(d, 1H, J = 9.0 Hz) 7.2-7.4(m, 1H) 6.93(d, 1H, J = 9.0 Hz) 6.8-7.1(m, 3H) 4.13(t, 2H, J = 6.9 Hz) 3.97(s, 3H) 3.72(q, 2H, J = 7.2 Hz) 2.95(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 6.9 Hz)	KBr 3134 2958 1670 1626 1535 1482 1286	FAB+ 427(M+H+) (100)
3-21  Colorless crystal / 120-121	CDCl ₃ , 300MHz 9.71(bt, 1H) 9.16(s, 1H) 8.67(s, 1H) 7.47(d, 1H, J = 8.8 Hz) 6.89(d, 1H, J = 7.9 Hz) 6.7-6.9(m, 3H) 5.61(s, 1H) 4.15(t, 2H, J = 6.9 Hz) 3.99(s, 3H) 3.90(s, 3H) 3.70(q, 2H, J = 7.4 Hz) 2.89(t, 2H, J = 7.4 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.96(t, 3H, J = 7.1 Hz)	3310 2952 1672 1625 1598 1529 1516 1260	FAB+ 455(M+H+) (100)
3-22  Colorless crystal / 139-140	CDCl ₃ , 300MHz 9.70(bt, 1H) 9.16(s, 1H) 8.85(s, 1H) 7.45(d, 1H, J = 8.9 Hz) 7.27(d, 2H, J = 8.4 Hz) 7.20(d, 2H, J = 8.4 Hz) 6.93(d, 1H, J = 8.9 Hz) 4.13(t, 2H, J = 6.9 Hz) 3.87(s, 3H) 3.70(q, 2H, J = 7.0 Hz) 2.92(t, 2H, J = 7.0 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.0 Hz)	KBr 3286 2960 1661 1613 1530 1496 1261	FAB+ 443(M+H+) (100)

Table 16




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-23  Colorless crystal / 127-128	CDCl ₃ , 300MHz 9.67(bt, 1H) 9.12(s, 1H) 8.86(s, 1H) 7.45(d, 1H, J = 8.7 Hz) 7.2-7.4(m, 5H) 6.93(d, 1H, J = 8.7 Hz) 4.13(t, 2H, J = 6.9 Hz) 3.97(s, 3H) 3.74(q, 2H, J = 7.2 Hz) 2.95(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.2 Hz)	KBr 3249 3139 2951 1661 1610 1284 1261 1116	FAB+ 409(M+H+) (100)
3-24  Colorless crystal / 133-134	CDCl ₃ , 300MHz 9.99(bt, 1H) 9.21(s, 1H) 8.91(s, 1H) 7.46(d, 1H, J = 8.9 Hz) 7.28(d, 2H, J = 7.9 Hz) 7.15(d, 2H, J = 7.9 Hz) 6.94(d, 1H, J = 8.9 Hz) 4.66(d, 2H, J = 5.7 Hz) 4.14(t, 2H, J = 6.9 Hz) 3.98(s, 3H) 2.34(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 3263 3417 1663 1618 1596 1587 1265 1262 1110	FAB+ 408(M+H+) (100)
3-25  Colorless crystal / 132-133	CDCl ₃ , 300MHz 10.01(bt, 1H) 9.13(s, 1H) 8.99(s, 1H) 7.47(d, 1H, J = 8.9 Hz) 7.3-7.4(m, 2H) 7.01(t, 2H, J = 8.7 Hz) 6.94(d, 1H, J = 8.9 Hz) 4.64(d, 2H, J = 5.8 Hz) 4.14(t, 2H, J = 6.9 Hz) 3.98(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.1 Hz)	KBr 3242 1664 1619 1537 1510 1263	FAB+ 413(M+H+) (100)

Table 17




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-26  Colorless crystal / 146-149	DMSO-d ₆ ,300MHz 11.42(brs,1H) 9.76(t,1H,J=5.4Hz) 8.75(s,1H) 8.47(d,2H,J=6.0Hz) 7.68(d,1H,J=9.3Hz) 7.30(d,2H,J=5.7Hz) 7.13(d,1H,J=9.0Hz) 3.95(m,5H) 3.64(q,2H,J=6.3Hz) 2.88(t,2H,J=6.9Hz) 1.78(m,2H) 0.94(t,3H,J=7.2Hz)	KBr 1671 1627 1537 1374 1265	(FAB+) 382(M+H+) (80)
3-27  Colorless crystal / 153-157	DMSO-d ₆ ,300MHz 11.42(brs1H) 9.75(t,1H,J=6.0Hz) 8.75(s,1H) 7.68(d,1H,J=8.7Hz) 7.31(m,2H) 7.12(m,3H) 3.95(m,5H) 3.58(q,2H,J=6.3Hz) 2.84(t,2H,J=6.9Hz) 1.78(m,2H) 0.94(t,3H,J=7.2Hz)	KBr 2961 1666 1624 1536 1219	(FAB+) 399(M+H+) (90)
3-28  Colorless crystal / 204-206	DMSO-d ₆ ,300MHz 11.40(brs,1H) 9.73(t,1H,5.4Hz) 9.16(s,1H) 8.74(s,1H) 7.68(d,1H,J=9.3Hz) 7.12(d,1H,J=9.3Hz) 7.06(d,2H,J=8.4Hz) 6.68(d,2H,J=8.4Hz) 3.95(m,5H) 3.52(q,2H,J=6.0Hz) 2.72(t,2H,J=6.9Hz) 1.78(m,2H) 0.94(t,3H,J=7.2Hz)	KBr 3236 1668 1611 1538 1261	(FAB+) 397(M+H+) (70)

Table 18




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-29  Colorless crystal / 140-143	DMSO-d ₆ , 300MHz 11.41(brs, 1H) 10.04(t, 1H, J=5.7Hz) 8.79(s, 1H) 7.70(d, 1H, J=8.7Hz) 7.14(d, 1H, J=8.7Hz) 6.86(m, 3H) 5.99(s, 2H) 4.47(d, 2H, J=5.7Hz) 3.96(m, 5H) 1.78(m, 2H) 0.94(t, 3H, J=7.5Hz)	KBr 2967 1664 1618 1535 1259	(FAB+) 411(M+H+) (80)
3-30  Colorless crystal / 136-138	DMSO-d ₆ , 300MHz 11.41(brs, 1H) 9.77(t, 1H, J=6.0Hz) 8.75(s, 1H) 7.68(d, 1H, J=8.7Hz) 7.26(m, 5H) 7.13(d, 1H, J=8.7Hz) 3.95(m, 5H) 3.59(q, 2H, J=5.7Hz) 2.85(t, 2H, J=7.2Hz) 1.78(m, 2H) 0.94(t, 3H, J=7.5Hz)	KBr 1666 1625 1537 1261 1113	(FAB+) 381(M+H+) (100)
3-31  Colorless crystal / 205-206	CDCl ₃ , 300MHz 9.65(bt, 1H) 9.26(s, 1H) 8.86(s, 1H) 7.47(d, 1H, J = 8.7 Hz) 7.2-7.3(m, 2H) 7.00(t, 2H, J = 8.4 Hz) 6.94(d, 1H, J = 8.7 Hz) 4.00(s, 3H) 3.98(s, 3H) 3.70(q, 2H, J = 7.2 Hz) 2.92(t, 2H, J = 7.2 Hz)	KBr 3650 3250 1664 1509 1221	FAB+ 371(M+H+) (100) 232(90)

Table 19




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-32  Colorless crystal / 257-258	DMSO-d ₆ , 300MHz 12.24(s, 1H) 9.89(bt, 1H) 7.45(s, 1H) 7.2-7.4(m, 2H) 7.11(t, 2H, J = 8.9 Hz) 6.90(s, 1H) 3.99(t, 2H, J = 6.6 Hz) 3.86(s, 3H) 3.55(q, 2H, J = 7.3 Hz) 2.83(t, 2H, J = 7.3 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.90(t, 3H, J = 7.0 Hz)	KBr 3448 1670 1560 1508 1266	FAB+ 427(M+H+) (100)
3-33  Colorless crystal / 252-253	CDCl ₃ , 300MHz 10.75(s, 1H) 10.00(bt, 1H) 8.88(s, 1H) 7.08(s, 1H) 6.7-6.9(m, 3H) 6.64(s, 1H) 5.93(s, 2H) 4.61(d, 2H, J = 5.8 Hz) 4.05(t, 2H, J = 6.8 Hz) 3.98(s, 3H) 1.8-2.0(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	3448 1671 1509 1266	FAB+ 439(M+H+) (100)
3-34  Colorless crystal / 188-189	DMSO-d ₆ , 300MHz 12.21(s, 1H) 9.98(bt, 1H) 8.71(s, 1H) 7.45(s, 1H) 6.90(s, 1H) 3.98(t, 2H, J = 6.9 Hz) 3.86(s, 3H) 3.5-3.7(m, 4H) 3.4-3.5(m, 2H) 2.6-2.7(m, 2H) 2.5-2.6(m, 4H) 1.6-1.9(m, 2H) 1.2-1.5(m, 4H) 0.93(t, 3H, J = 7.0 Hz)	KBr 3448 1676 1244	FAB+ 418(M+H+) (100)

Table 20




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-35  Colorless crystal / 158-160	DMSO-d ₆ , 300MHz 11.54(brs, 1H) 10.05(t, 1H, J=5.9Hz) 8.79(s, 1H) 7.70(d, 1H, J=8.8Hz) 7.14(d, 1H, J=8.8Hz) 6.87(m, 3H) 5.99(s, 2H) 4.47(d, 2H, J=5.9Hz) 4.07(q, 2H, J=7.0Hz) 3.93(s, 3H) 1.32(t, 3H, J=7.0Hz)	KBr 3168 1673 1619 1536 1257 1106	(FAB+) 397(M+H+) (80)
3-36  Colorless crystal / 136-137	CDCl ₃ , 300MHz 9.90(bt, 1H) 8.74(s, 1H) 7.46(d, 1H, J = 8.7 Hz) 7.2-7.3(m, 2H) 6.9-7.1(m, 2H) 6.97(d, 1H, J = 8.7 Hz) 4.11(s, 3H) 4.09(s, 3H) 3.86(t, 2H, J = 6.8 Hz) 3.69(q, 2H, J = 7.3 Hz) 2.92(t, 2H, J = 7.3 Hz) 1.7-1.9(m, 2H) 1.3-1.6(m, 4H) 0.94(t, 3H, J = 7.1 Hz)	3231 2953 1673 1612 1535 1530 1270 1219	FAB+ 441(M+H+) (100) 302(90)
3-37  Colorless crystal / 127-128	CDCl ₃ , 300MHz 9.97(bt, 1H) 8.74(s, 1H) 8.52(d, 2H, J = 5.9 Hz) 7.48(d, 1H, J = 8.8 Hz) 7.21(d, 2H, J = 5.9 Hz) 6.99(d, 1H, J = 8.8 Hz) 4.00(s, 3H) 3.98(s, 3H) 3.87(t, 2H, J = 6.8 Hz) 3.75(q, 2H, J = 7.2 Hz) 2.96(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.1 Hz)	KBr 3448 2955 1672 1611 1579 1334 1452 1270	FAB+ 424(M+H+) (100)

Table 21




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-38  Colorless crystal / 102-103	CDCl ₃ , 300MHz 9.96(bt, 1H) 8.74(s, 1H) 7.46(d, 1H, J = 8.7 Hz) 6.97(d, 1H, J = 8.7 Hz) 4.02(s, 3H) 3.98(s, 3H) 3.85(t, 2H, J = 6.6 Hz) 3.7-3.8(m, 4H) 3.61(q, 2H, J = 6.6 Hz) 2.62(t, 2H, J = 6.6 Hz) 2.5-2.6(m, 4H) 1.8-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.2 Hz)	KBr 3448 2948 1672 1612 1581 1560 1535 1452 1269	FAB+ 432(M+H+) (100)
3-39  Colorless crystal / 146-147	CDCl ₃ , 300MHz 10.38(bt, 1H) 8.78(s, 1H) 8.55(d, 2H, J = 6.8 Hz) 7.49(d, 1H, J = 8.7 Hz) 7.28(d, 2H, J = 6.8) 7.00(d, 1H, J = 8.7 Hz) 4.69(t, 2H, J = 6.0 Hz) 4.08(s, 3H) 3.99(s, 3H) 3.89(t, 2H, J = 6.8 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 7.1 Hz)	KBr 3264 2957 1668 1609 1579 1526 1286	FAB+ 410(M+H+) (100)
3-40  Colorless crystal / 132-133	CDCl ₃ , 300MHz 10.22(bt, 1H) 8.78(s, 1H) 7.48(d, 1H, J = 8.7 Hz) 7.3-7.4(m, 2H) 7.02(d, 1H, J = 8.7 Hz) 6.9-7.0(m, 3H) 4.64(d, 2H, J = 5.8 Hz) 4.00(s, 3H) 3.98(s, 3H) 3.87(t, 2H, J = 6.9 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 6.9 Hz)	KBr 3448 1688 1610 1579 1560 1528 1288	FAB+ 427(M+H+) (100)

Table 22




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-41  Colorless crystal / 180-181	CDCl ₃ ,300MHz 10.01(bt, 1H) 8.75(s, 1H) 7.46(d, 1H, J = 8.8 Hz) 7.08(d, 2H, J = 8.4 Hz) 6.73(d, 1H, J = 8.8 Hz) 6.79(d, 2H, J = 8.4 Hz) 6.90(bs, 1H) 4.00(s, 3H) 3.98(s, 3H) 3.86(t, 2H, J = 6.8 Hz) 3.68(q, 2H, J = 7.2 Hz) 2.86(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.1 Hz)	KBr 3246 2939 1672 1606 1536 1514 1500 1270	FAB+ 439(M+H+) (100)
3-42  Colorless crystal / 107-108	CDCl ₃ ,300MHz 10.15(bt, 1H) 8.77(s, 1H) 7.47(d, 1H, J = 8.8 Hz) 6.98(d, 1H, J = 8.8 Hz) 6.87(d, 1H, J = 1.5 Hz) 6.84(dd, 1H, J = 7.7, 1.5 Hz) 6.75(d, 1H, J = 7.7 Hz) 5.92(s, 2H) 4.57(d, 2H, J = 5.5 Hz) 4.00(s, 3H) 3.98(s, 3H) 3.87(t, 2H, J = 7.0 Hz) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.94(t, 3H, J = 7.0 Hz)	KBr 2930 1268 800	FAB+ 453(M+H+) (100) 302(100)
3-43  Colorless crystal / 147-148	CDCl ₃ ,300MHz 10.00(bt, 1H) 8.77(s, 1H) 7.2-7.3(m, 2H) 7.10(s, 1H) 6.99(t, 2H, J = 8.6 Hz) 6.77(s, 1H) 4.07(t, 2H, J = 7.0 Hz) 4.02(s, 3H) 3.77(s, 3H) 3.70(q, 2H, J = 7.7 Hz) 2.92(t, 2H, J = 7.7 Hz) 1.7-2.0(m, 2H) 1.3-1.6(m, 4H) 0.95(t, 3H, J = 6.9 Hz)	KBr 2935 1672 1510 1426 1260	FAB+ 442(M+H+) (100) 302(100)

Table 23




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
3-44  Colorless crystal / 174-175	CDCl ₃ , 300MHz 10.06(bt, 1H) 8.77(s, 1H) 7.10(s, 1H) 6.77(s, 1H) 4.07(t, 2H, J = 7.0 Hz) 4.03(s, 3H) 3.78(s, 3H) 3.6-3.7(m, 4H) 3.62(q, 2H, J = 6.6 Hz) 2.63(t, 2H, J = 6.6 Hz) 2.5-2.6(m, 4H) 1.8-2.0(m, 2H) 1.3-1.6(m, 4H) 0.95(t, 3H, J = 7.0 Hz)	KBr 2934 1672 1609 1543 1430 1397 1260 1171 1120	FAB+ 432(M+H+) (100) 302(100)
3-45  Colorless crystal / 176-177	CDCl ₃ , 300MHz 10.24(t, 1H, J = 5.9 Hz) 8.81(s, 1H) 7.11(s, 1H) 6.88(s, 1H) 6.84(d, 1H, J = 8.1 Hz) 6.78(s, 1H) 6.75(d, 1H, J = 8.1 Hz) 5.92(s, 2H) 4.58(d, 2H, J = 5.9 Hz) 4.07(t, 2H, J = 7.0 Hz) 4.02(s, 3H) 3.76(s, 3H) 1.8-2.0(m, 2H) 1.3-1.6(m, 4H) 0.95(t, 3H, J = 7.0 Hz)	KBr 3448 1671 1542 1251	FAB+ 453(M+H+) (100) 302(100)
3-46  Colorless crystal / 106-107	CDCl ₃ , 300MHz 9.67(bt, 1H) 9.15(s, 1H) 8.85(s, 1H) 7.42(d, 1H, J = 8.7 Hz) 7.2-7.3(m, 2H) 7.00(t, 2H, J = 8.4 Hz) 6.91(d, 1H, J = 8.7 Hz) 4.15(t, 2H, J = 7.2 Hz) 4.12(t, 2H, J = 6.6 Hz) 3.70(q, 2H, J = 7.2 Hz) 2.92(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 4H) 1.3-1.6(m, 8H) 0.95(t, 3H, J = 7.2 Hz) 0.95(t, 3H, J = 7.2 Hz)	KBr 3286 1654 1520 1500	FAB+ 483(M+H+) (100) 344(50)

Table 24




Ex.comp./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR m ⁻¹	MS
3-47  Colorless crystal / 241~242	DMSO-d ₆ , 300MHz 11.22 (bs, 1H) 10.10 (t, 1H, J = 5.8 Hz) 9.66 (s, 1H) 8.76 (s, 1H) 7.45 (d, 1H, J = 8.8 Hz) 7.14 (d, 1H, J = 8.8 Hz) 6.91 (s, 1H) 6.87 (d, 1H, J = 7.9 Hz) 6.82 (d, 1H, J = 7.9 Hz) 5.99 (s, 2H) 4.45 (d, 2H, J = 5.8 Hz) 3.91 (s, 3H)	KBr 3161 1655 1268 1113 1039 934 802 521	LCQ(+) 369 [M+H] (100)
3-48  Colorless crystal / 124~125	DMSO-d ₆ , 300MHz 11.42 (s, 1H) 9.96 (t, 1H, J = 5.7Hz) 8.86 (s, 1H) 8.79 (s, 1H) 8.75 (s, 1H) 7.67 (d, 1H, J = 9.0 Hz) 7.14 (d, 1H, J = 9.0 Hz) 6.73 (d, 1H, J = 3.6 Hz) 6.68 (d, 1H, J = 8.1 Hz) 6.59 (dd, 1H, J = 8.1, 3.6 Hz) 4.38 (d, 2H, J = 5.7 Hz) 3.99 (t, 2H, J = 6.9 Hz) 3.93 (s, 3H) 1.7-1.8 (m, 2H) 1.3-1.4 (m, 4H) 0.89 (t, 3H, J = 7.2 Hz)	KBr 3246 1672 1626 1536 1260 1109	FAB+ 427 [M+H] (100)
3-49  Colorless crystal / 145~146	DMSO-d ₆ , 300MHz 11.0 (bs, 1H) 10.2 (bs, 1H) 10.17 (bt, 1H) 8.76 (s, 1H) 7.66 (d, 1H, J = 8.9 Hz) 7.14 (d, 1H, J = 8.9 Hz) 6.93 (s, 1H) 6.74 (s, 2H) 4.45 (d, 2H, J = 5.5 Hz) 3.98 (t, 2H, J = 6.8 Hz) 3.91 (s, 3H) 3.75 (s, 3H) 1.7-1.9 (m, 2H) 1.3-1.5 (m, 4H) 0.88 (t, 3H, J = 7.0 Hz)	KBr 3350 3193 2954 1920 1668 1627 1528 1464	LCQ(-) 439 [M-H] (100)

Table 25




Ex.comp./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR m ⁻¹	MS
3-50  Colorless crystal /	DMSO-d ₆ , 300MHz 11.45, 11.39(s, 1H) 9.9-10.1(t, 1H) 8.79, 8.79(s, 1H) 6.6-6.8(bs, 1H) 7.64(d, 1H, J = 8.9Hz) 7.13(d, 1H, J = 8.9Hz) 6.7-7.1(m, 3H) 5.63, 5.57(bs, 1H) 5.22(bs, 1H), 4.8-4.9(m, 1H), 4.44(bd, 2H) 3.98(t, 2H, J = 6.9Hz) 3.93(s, 3H) 3.82(d, 1H, J = 9.5 Hz) 3.3-3.5(m, 4H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.89(t, 3H, 7.2Hz)	KBr 3396 1666	FAB+ 603 [M+H] (100)
3-51  Colorless crystal / 209~210	DMSO-d ₆ , 300MHz 11.8(bs, 1H) 11.5(bs, 1H) 10.04(bt, 1H) 8.79(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.91(s, 1H) 6.86(d, 1H, J = 7.9 Hz) 6.82(d, 1H, J = 7.9 Hz) 5.98(s, 2H) 4.46(d, 2H, J = 5.9 Hz) 3.99(t, 2H, J = 6.5 Hz) 3.93(s, 3H) 2.27(t, 2H, J = 7.2 Hz) 1.7-1.9(m, 2H) 1.6-1.7(m, 2H)	KBr 3300 1730	LCQ(+) 469 [M+H] (100)
3-52  Colorless crystal / 189~190	DMSO-d ₆ , 300MHz 12.00(bs, 1H) 11.50(bs, 1H) 10.00(t, 1H, J = 5.7 Hz) 8.88(bs, 1H) 8.80(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.93(s, 1H) 6.74(s, 2H) 4.45(d, 2H, J = 5.7 Hz) 3.98(t, 2H, J = 6.6 Hz) 3.96(s, 3H) 3.75(s, 3H) 2.2-2.4(m, 2H) 1.7-1.8(m, 2H) 1.6-1.7(m, 2H)	KBr 3449 1707 1686 1626 1545 1499 1263	LCQ(+) 471 [M+H] (100)

Table 26




Ex.comp./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm^{-1}	MS
3-53  Colorless crystal / 96~97	DMSO-d ₆ , 300MHz 11.45(s, 1H) 10.04(t, 1H, J = 5.9 Hz) 8.79(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.91(s, 1H) 6.87(d, 1H, J = 7.9 Hz) 6.82(d, 1H, J = 7.9 Hz) 5.99(s, 2H) 4.47(d, 2H, J = 5.9 Hz) 4.36(t, 2H, J = 5.1 Hz) 3.98(t, 2H, J = 6.9 Hz) 3.93(s, 3H) 3.3-3.4(m, 2H) 1.7-1.8(m, 2H) 1.4-1.5(m, 4H)	KBr 3300 2900 1550	LCQ(+) 455 [M+H] (100)
3-54  Colorless amorphous solid /	DMSO-d ₆ , 300MHz 11.46(s, 1H) 10.00(t, 1H, J = 5.7 Hz) 8.88(s, 1H) 8.80(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.93(s, 1H) 6.73(s, 2H) 4.45(d, 1H, J = 5.7 Hz) 4.36(t, 1H, J = 5.3 Hz) 3.98(t, 2H, J = 6.8 Hz) 3.93(s, 3H) 3.74(s, 3H) 3.3-3.5(m, 2H) 1.7-1.9(m, 2H) 1.4-1.6(m, 4H)	KBr 3392 1554 1260 1110 1038 801	LCQ(+) 457 [M+H] (100)
3-55  Colorless crystal / 128~129	DMSO-d ₆ , 300MHz 11.47(s, 1H), 10.04(t, 1H, J = 6.0 Hz) 8.79(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.91(d, 1H, J = 1.4 Hz) 6.87(d, 1H, J = 7.9 Hz) 6.82(dd, 1H, J = 7.9, 1.4 Hz), 5.99(s, 2H) 4.47(d, 2H, J = 6.0 Hz) 4.43(d, 1H, J = 4.7 Hz) 3.99(t, 2H, J = 6.9 Hz) 3.93(s, 3H) 3.6-3.7(m, 1H) 1.7-1.9(m, 2H) 1.4-1.5(m, 2H) 1.07(d, 3H, J = 6.2 Hz)	KBr 3397 2965 1672 1623 1544 1501 1260	LCQ(+) 455 [M+H] (100)

Table 27




Ex.comp./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR m ⁻¹	MS
3-56  Colorless crystal / 153~154	DMSO-d ₆ , 300MHz 11.53(s, 1H) 10.04(t, 1H, J = 5.9 Hz) 8.79(s, 1H) 7.70(d, 1H, J = 8.9 Hz) 7.14(d, 1H, J = 8.9 Hz) 6.91(s, 1H) 6.87(d, 1H, J = 7.9 Hz) 6.82(d, 1H, J = 7.9 Hz) 5.99(s, 2H) 4.47(d, 2H, J = 5.9 Hz) 3.96(t, 2H, J = 6.5 Hz) 3.92(s, 3H) 2.64(t, 2H, J = 7.2Hz) 2.13(s, 3H) 1.9-2.0(m, 2H)	KBr 3290 2895 1707 1672 1621 1538 1440 1371	LCQ(+) 453 [M+H] ⁺ (100)
3-57  Colorless crystal / 118~119	DMSO-d ₆ , 400MHz 11.55(s, 1H) 10.02(bt, 1H) 8.78(s, 1H) 7.69(d, 1H, J = 8.9 Hz) 7.15(d, 1H, J = 8.9 Hz) 6.90(s, 1H) 6.87(d, 1H, J = 8.0 Hz) 6.82(d, 1H, J = 8.0 Hz) 5.98(s, 2H) 4.82(bs, 1H) 4.46(d, 2H, J = 5.8Hz) 4.0-4.2(2H), 3.93(s, 3H), 3.58(bs, 1H) 1.8-2.0(m, 1H) 1.6-1.8(m, 1H) 1.4-1.5(m, 2H) 0.88(t, 3H, J = 7.4 Hz)	KBr 3438 1626 1110 1802	LCQ(+) 455 [M+H] ⁺ (100)
3-58  Colorless crystal / 148~149	DMSO-d ₆ , 300MHz 11.31(s, 1H) 10.01(bt, 1H) 8.81(s, 1H) 7.74(d, 1H, J = 8.7 Hz) 7.15(d, 1H, J = 8.9 Hz) 6.92(d, 1H, J = 0.9 Hz) 6.85(d, 1H, J = 7.8 Hz) 6.82(dd, 1H, J = 7.8, 0.9 Hz) 5.99(s, 2H) 4.46(d, 2H, J = 5.7 Hz) 4.15(t, 2H, J = 6.0 Hz) 3.93(s, 3H) 3.01(t, 2H, J = 5.4Hz) 2.55(q, 2H, 6.9 Hz) 0.97(t, 3H, J = 6.9 Hz)	KBr 3448 1676 1534 1259 1120 805	LCQ(+) 453 [M+H] ⁺ (100)

Table 28



Ex.comp./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR m ⁻¹	MS
3-59  Colorless crystal / 185~186	DMSO-d ₆ , 300MHz 11.47(s, 1H) 10.00(t, 1H, J = 6.0 Hz) 8.81(s, 1H) 7.73(d, 1H, J = 8.7 Hz) 7.15(d, 1H, J = 8.7 Hz) 6.90(s, 1H) 6.87(d, 1H, J = 7.8 Hz) 6.81(d, 1H, J = 7.8 Hz) 5.99(s, 2H) 5.65(bd, 1H) 4.46(d, 2H, J = 6.0 Hz) 4.07(dd, 1H, J = 9.9, 2.1 Hz), 3.94(s, 3H) 3.80(bs, 1H) 3.6-3.7(m, 1H) 1.3-1.5(m, 4H) 0.89(t, 3H, J = 6.6 Hz)	KBr 3404 1666 1545 1264	LCQ(+) 455 [M+H] (100)
3-60  Colorless crystal / 198~199	CDCl ₃ , 300MHz 9.36(bs, 1H) 9.23(s, 1H) 8.88(s, 1H) 7.45(d, 1H, 9.0 Hz) 6.94(d, 1H, 9.0 Hz) 5.86(bs, 1H) 4.14(t, 2H, J = 7.2 Hz) 3.98(s, 3H) 1.7-1.9(m, 2H) 1.3-1.5(m, 4H) 0.95(t, 3H, J = 6.9 Hz)	KBr 3329 3163 2956 1687 1500 1370 1300	FAB+ 305 [M+H] (100)

Table 29




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
4-1  Pale yellow crystal / 272-273	DMSO-d ₆ , 300MHz 11.06(s, 1H) 10.30(s, 1H) 9.80(bt, 1H) 9.25(s, 1H) 8.65(s, 1H) 7.29(d, 1H, J = 8.6 Hz) 7.2-7.4(m, 2H) 7.1-7.2(m, 2H) 6.83(d, 1H, J = 8.6 Hz) 3.55(q, 2H, J = 7.2 Hz) 2.83(t, 2H, J = 7.2 Hz)	KBr 3236 1668 1510 1357 1220	FAB+ 343[M-H+] (100)
5-1  Colorless crystal / 99-100	CDCl ₃ , 300MHz 9.10 (br s, 1H) 7.65 (s, 1H) 7.26 (d, J=8.0 Hz, 1H) 6.86 (d, J=8.0 Hz, 1H) 4.65 (s, 2H) 4.13 (t, J=6.6 Hz, 2H) 3.95 (s, 3H) 1.80 (m, 2H) 1.50 (m, 2H) 0.99 (t, J=7.3 Hz, 3H)	KBr 3176 1651 1610 1509 1281 1111 1063	(fab+, NBA) 278 260 (fab-, NBA) 276 199
5-2  Colorless crystal / 154-157	CDCl ₃ , 300MHz 9.18(brs, 1H) 7.65(s, 1H) 7.26(d, 1H, J=8.4Hz) 6.87(d, 1H, J=8.7Hz) 4.65(s, 2H) 4.22(q, 2H, 6.9Hz) 3.95(s, 3H) 3.36(brs, 1H) 1.42(t, 3H, J=6.9Hz)	KBr 3434 1644 1510 1379 1283	

Table 30




Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
6-1  Colorless crystal / 164-165	CDCl ₃ , 300MHz 9.05 (br s, 1H) 7.83 (s, 1H) 7.33 (dd, J=4.6, 8.9 Hz, 2H) 7.26 (d, J=8.8 Hz, 1H) 6.99 (t, J=8.8 Hz, 2H) 6.86 (br s, 1H) 6.85 (d, J=8.8 Hz, 1H) 5.21 (s, 2H) 4.11 (t, J=6.8 Hz, 2H) 3.94 (s, 3H) 1.79 (m, 2H) 1.50 (m, 2H) 0.99 (t, J=7.7 Hz, 3H)	KBr 3305 1706 1656 1611 1537 1509 1262 1219 1112 1077	(fab+, NBA) 415 260 (fab-, NBA) 413 276 199
6-2  Colorless crystal / 205(dec.)	DMSO-d ₆ , 300MHz 11.08 (brs, 1H) 10.27 (s, 1H) 8.39 (d, 2H, J=6.2 Hz) 7.94 (s, 1H) 7.44 (m, 3H) 7.02 (d, 1H, J=8.8 Hz) 5.06 (s, 2H) 4.05 (q, 2H, J=7.0 Hz) 3.89 (s, 3H) 1.31 (t, 3H, J=7.0 Hz)	KBr 2978 1745 1659 1609 1508 1251 1209	(FAB+) 370 (M+H+) (50)
7-1  Colorless crystal / 134-136	CDCl ₃ , 300MHz 9.03 (brs, 1H) 7.75 (s, 1H) 7.26 (d, 1H, J=8.8 Hz) 6.84 (d, 1H, J=8.8 Hz) 4.20 (q, 2H, J=7.0 Hz) 3.94 (s, 3H) 3.47 (s, 2H) 2.35 (s, 6H) 1.41 (t, 3H, J=7.0 Hz)	KBr 2773 1644 1605 1285 1109	FAB+ 277 (M+H+) (100)

Table 31





Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
7-2  Colorless crystal /	CDCl ₃ ,300MHz 9.00 (brs, 1H) 7.65 (s, 1H) 7.24 (d, J=8.7Hz) 6.83 (d, J=8.7 Hz, 1H) 4.12 (t, J=6.9 Hz, 2H) 3.93 (s, 3H) 3.82 (s, 2H) 1.80 (m, 2H) 1.80 (brs, 2H) 1.50 (m, 2H) 0.99 (t, J=7.3 Hz, 3H)		(fab+, NBA) 278 260 (fab-, NBA) 276 199
7-3  Colorless crystal / 166-169	CDCl ₃ ,300MHz 9.03(brs,1H) 7.78(brs,1H) 7.27(d,1H,J=8.8Hz) 6.85(d,1H,J=8.8Hz) 4.20(q,2H,J=7.0Hz) 3.94(s,3H) 3.78(t,4H,4.8Hz) 3.56(brs,2H) 1.41(t,3H,J=7.0Hz)	KBr 2968 1656 1611 1280 1115	FAB+ 319(M+H+) (100)
8-1  Colorless crystal / 207-209	DMSO-d ₆ ,300MHz 8.75 (br s, 1H) 7.70 (s, 1H) 7.43-7.38 (m, 3H) 7.05 (t, J=8.9 Hz, 2H) 6.98 (d, J=8.9 Hz, 1H) 6.54 (br t, J=5.8 Hz, 1H) 4.04 (d, J=5.8 Hz, 2H) 3.98 (t, J=6.8 Hz, 2H) 3.87 (s, 3H) 1.75 (m, 2H) 1.41 (m, 2H) 0.93 (t, J=7.3 Hz, 3H)	KBr 3313 1659 1611 1577 1508 1281 1216 1113	

Table 32

Ex.No./structure/character/mp (°C)	¹ H NMR (δ) ppm	IR cm ⁻¹	MS
8-2  Colorless crystal / 199-200	DMSO-d ₆ , 300MHz 10.72 (br s, 1H) 9.24 (br s, 1H) 8.27 (t, J=5.9 Hz, 1H) 7.41 (s, 1H) 7.22 (d, J=8.8 Hz, 1H) 7.11 (d, J=8.4 Hz, 2H) 6.97 (d, J=8.8 Hz, 1H) 6.71 (d, J=8.4 Hz, 2H) 4.07 (d, J=5.5 Hz, 2H) 3.97 (t, J=6.6 Hz, 2H) 3.87 (s, 3H) 3.38 (s, 2H) 1.74 (m, 2H) 1.41 (m, 2H) 0.92 (t, J=7.3 Hz, 3H)	KBr 3387 1648 1609 1509 1278 1260 1108	

Pharmacological Experiment

[1] Binding assay (*in vitro*)

[0203] Membrane fractions of human central type cannabinoid receptor (human CB1-CHO, hereinafter called hCB1) and human peripheral type cannabinoid receptor (human CB2-CHO, hereinafter called hCB2) expressed in CHO cells were used as samples. The reference material (hCB1: 20 µg/ml, hCB2: 5 µg/ml), labeled ligand ([³H]Win55212-2, 2 nM) and unlabeled ligand Win55212-2 or test compound were added to a 24-well round bottom plate, and the plate was incubated at 30°C for 90 minutes. 50 mM Tris-HBSS containing 0.2% BSA was used as an assay buffer. After incubation, the mixture was filtered (Packard, Unifilter 24GF/B) and dried, and then a scintillation solution (Packard, Microsint-20) was added thereto to determine the radioactivity of the sample (Packard, Top count A9912V). Nonspecific binding was determined by the addition of excessive amount of Win55212-2 (10 µM), and specific binding was calculated by subtracting the nonspecific binding from total binding determined by the addition of the labeled ligand only. The test compound was dissolved in DMSO to make the final concentration in DMSO to be 0.1%. IC₅₀ value was determined from the proportion of the bound test compound in the specific binding, and K_i value of the test compound was calculated from the IC₅₀ value and K_d value of [³H]Win55212-2. K_i value for the central type receptors/K_i value for peripheral type receptors (C/S) was also calculated as an indicator of the selectivity of the test compound for peripheral type receptors. Results are shown in Tables 33 to 36.

Table 33

Example	K _i value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
1-2	3671	0.014	262202
1-4	1627	4.49	362
2-1	4330	8.90	487
2-2	3247	00.77	42172
2-4	905	0.032	28273
2-5	434	0.20	2170
2-6	770	0.13	5923
2-7	381	0.42	908

EP 1 142 877 A1

Table 33 (continued)

Example	Ki value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
3-1	49	0.13±0.05	372
3-5	3436	0.087	39497
3-6	609	0.020	30472
3-7	72	0.48	150
3-8	81	0.019	4275

Table 34

Example	Ki value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
3-14	131	0.33	396
3-15	249	0.016	15580
3-16	208	0.010	20805
3-17	199	0.23	867
3-18	1751	7.10	247
3-19	49	0.18	272
3-20	78	0.41	191
3-21	162	0.97	167
3-22	135	0.20	675
3-23	39	0.20	194
3-25	444	1.29	344
3-26	648	0.23	2817
3-27	336	0.021	15990

Table 35

Example	Ki value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
3-30	168	0.090	1862
3-31	159	0.16	995
3-32	2398	0.036	66604
3-33	273	1.06	258
3-34	172	0.011	15672
3-35	409	0.053	7713
3-36	183	0.021	8695
3-37	78	0.75	104
3-38	935	0.085	11002
3-40	703	2.13	330
3-41	62	0.35	176

Table 35 (continued)

Example	Ki value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
3-42	315	0.22	1430
3-43	864	0.043	20093

Table 36

Example	Ki value (nM)		
	Central type receptor (C)	Peripheral type receptor (S)	C/S
3-44	193	0.18	1075
3-45	381	0.032	11902
3-46	228	0.026	8776
5-1	686	0.45	1525
5-2	122	0.055	2219
6-1	703	0.045	15612
6-2	437	0.034	12866
7-1	3643	0.17	21432
7-3	666	0.062	10735
8-1	440	0.44	1000
8-2	727	0.27	2692

[2] Carrageenin-induced paw edema model (*in vivo*)

[0204] C57BL/6J mice (6 to 8 weeks of age) were used. The volume of the right hind paw before administration was measured (Unicom, Prethysumometer TK-101), and 2 hours later, the test compound dissolved in olive oil was administered orally at 10 ml/kg. After one hour of the administration, 50 μ l of 1% solution of carrageenin in normal saline was administered intradermally at foot pad of the right hind paw. Three hours later, the volume of the right hind paw was measured and compared with the initial volume. Results are shown in Table 37.

Table 37

Example	ED ₅₀ (mg/kg : p.o.)	Example	ED ₅₀ (mg/kg : p.o.)
3-5	<0.10	3-26	1.32
3-9	0.58	3-27	0.53
3-11	0.40	3-30	0.49
3-15	0.41	3-37	0.63
3-17	1.39	3-38	2.24
3-22	0.76	3-40	1.13

[3] The compounds of the invention were shown to significantly inhibit inflammation and hemorrhage in the pancreas, in the experiments using pancreatitis model induced by taurocholic acid in rats.

[0205] An example of formulation is shown below, but is not intended to limit the present invention.

Example of formulation

[0206]

- (a) The compound of Example 1-1: 10 g
- (b) Lactose: 50 g
- (c) Maize starch: 15 g
- (d) Carboxymethylcellulose sodium: 44 g
- (e) Magnesium stearate: 1 g


[0207] All of (a), (b) and (c), and 30 g of (d) were kneaded with water, and the mixture was dried in vacuo and then granulated. To this granular powder, 14 g of (d) and 1 g of (e) were mixed, and the mixture was tableted by a tablet maker to manufacture 1000 tablets containing 10 mg of (a) in a tablet.

Industrial Applicability

[0208] The compound [I] of the present invention and its pharmaceutically acceptable salts selectively act on cannabinoid receptors, particularly on peripheral type cannabinoid receptors, and have fewer side effects on the central nervous system, having great immunomodulating action, anti-inflammatory action and antiallergic action. Therefore, these compounds are useful as cannabinoid receptor (particularly peripheral cannabinoid receptors) regulator, immunomodulators, therapeutic agent for autoimmune diseases, anti-inflammatory agents, and antiallergic agents.

Claims

1. A cannabinoid receptor modulator comprising, as an active ingredient, a 2-oxoquinoline compound represented by the following general formula [I] or its pharmaceutically acceptable salt:



[1]

where W means -O-, -S(O)_t-, -CR³R⁴-, -NR⁵-, -NR⁵CO-, -CONR⁵-, -COO- or -OCO- (where R³ and R⁴ may be identical or different and represent hydrogen atom or alkyl; R⁵ represents hydrogen atom or alkyl; and t indicates an integer, 0, 1 or 2);


R¹ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; each group of R¹, except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, carboxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkylsulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R² represents hydrogen atom, alkyl, -OR⁶ (where R⁶ represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl), -NR⁷R⁸ (where R⁷ and R⁸ may be identical or different and represent hydrogen atom, alkyl, alkenyl, alkynyl, acyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; or R⁷ and R⁸, together with the adjacent nitrogen atom, can form a heteroaryl); or -(CH₂)_u-S(O)_{u'}R⁹ (where R⁹ represents hydrogen atom, alkyl, alkenyl or alkynyl, each of u and u' independently represents an integer, 0, 1 or 2); each group of R², except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkyl sulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R^a represents hydrogen atom or alkyl;


X represents -COOR^b, -CONH₂, -CONR^c-(Alk^a)_r-R, -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R, -(CH₂)_q-NR^eC(=Z)-(NR^f)_w-(Alk^c)_v-R, -(CH₂)_p-OH or -(CH₂)_q-NR^eR^{e'}

(where each of R^b, R^c, R^d, and R^f independently represents hydrogen atom or alkyl; each of R^e and R^{e'} independently represents hydrogen atom or alkyl; or R^e and R^{e'}, together with the adjacent nitrogen atom, can form a heteroaryl; each of Alk^a, Alk^b and Alk^c independently represents alkylene or alkenylene; each of the alkylene and alkenylene may be substituted or unsubstituted with hydroxy, carboxy, alkoxycarbonyl, alkyl (the alkyl may be substituted or unsubstituted with hydroxy, alkoxy or alkylthio) or -CONR¹⁰R¹¹ (where R¹⁰ and R¹¹ may be identical or different and represent hydrogen atom or alkyl; or R¹⁰ and R¹¹, together with the adjacent nitrogen atom, can form a heteroaryl); R represents aryl, heteroaryl, cycloalkyl, benzene-condensed cycloalkyl or




where A and B independently represent oxygen atom, nitrogen atom or sulfur atom; k indicates an integer of 1-3; each of the aryl and heteroaryl may be substituted or unsubstituted with an alkyl substituted or unsubstituted with hydroxy or may be substituted or unsubstituted with a hydroxy, alkoxy, alkenyloxy, acyl, acyloxy, halogen atom, nitro, amino, sulfonamide, alkylamino, aralkyloxy, pyridyl, piperidino, carboxy, alkoxycarbonyl, acylamino, aminocarbonyl, cyano or glucuronic acid residue; the cycloalkyl may be substituted or unsubstituted with a hydroxy, alkoxy or =O; the benzene-condensed cycloalkyl may be substituted or unsubstituted with a hydroxy or alkoxy; each of r, s, v and w independently represents 0 or 1; each of Y and Z independently represents a nitrogen atom, oxygen atom or sulfur atom; and each of p and q independently represents an integer of 1-4).

2. A cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of claim 1 or its pharmaceutically acceptable salt, wherein W represents -O-; R¹ is hydrogen atom or alkyl (the alkyl is as described above); R² represents -OR⁶ (R⁶ is as described above); and R is aryl, heteroaryl or



where aryl, heteroaryl, and each symbol in this formula are as defined above.

3. A 2-oxoquinoline compound as represented by the following general formula [I'] or its pharmaceutically acceptable salt:



where W means $-O-$, $-S(O)_t-$, $-CR^3R^4-$, $-NR^5-$, $-NR^5CO-$, $-CONR^5-$, $-COO-$ or $-OCO-$ (where R^3 and R^4 may be identical or different and represent hydrogen atom or alkyl; R^5 represents hydrogen atom or alkyl; and t indicates an integer, 0, 1 or 2);

R^1 represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; each group of R^1 , except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, carboxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkylsulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;

R^2 represents hydrogen atom, alkyl, $-OR^6$ (where R^6 represents hydrogen atom, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl), $-NR^7R^8$ (where R^7 and R^8 may be identical or different and represent hydrogen atom, alkyl, alkenyl, alkynyl, acyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl; or R^7 and R^8 , together with the adjacent nitrogen atom, can form a heteroaryl); or $-(CH_2)_u-S(O)_vR^9$ (where R^9 represents hydrogen atom, alkyl, alkenyl or alkynyl, each of u and v independently represents an integer, 0, 1 or 2); each group of R^2 , except hydrogen atom, may be substituted or unsubstituted with alkylamino, amino, hydroxy, alkoxy, alkoxycarbonyl, acyl, acyloxy, acylthio, mercapto, alkylthio, alkylsulfinyl or alkyl sulfonyl; each group, except hydrogen atom and alkyl, may be substituted or unsubstituted with alkyl;


R^a represents hydrogen atom or alkyl;

X' represents $-CONR^c-(Alk^a)_r-R$, $-(CH_2)_p-OC(=Y)-NR^d-(Alk^b)_s-R$ or $-(CH_2)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R$, where each of R^c , R^d , R^e and R^f independently represents hydrogen atom or alkyl;

each of Alk^a , Alk^b and Alk^c independently represents alkylene or alkenylene; each of the alkylene and alkenylene may be substituted or unsubstituted with hydroxy, carboxy, alkoxycarbonyl, alkyl (the alkyl may be substituted or unsubstituted with a hydroxy, alkoxy or alkylthio);

or $-CONR^{10}R^{11}$ (where R^{10} and R^{11} may be identical or different and represent hydrogen atom or alkyl; or R^{10} and R^{11} , together with the adjacent nitrogen atom, can form a heteroaryl);

R represents aryl, heteroaryl, cycloalkyl, benzene-condensed cycloalkyl or




where A and B independently represent oxygen atom, nitrogen atom or sulfur atom; k indicates an integer of 1-3; each of the aryl and heteroaryl may be substituted or unsubstituted with an alkyl substituted or unsubstituted with hydroxy or may be substituted or unsubstituted with a hydroxy, alkoxy, alkenyloxy, acyl, acyloxy, halogen atom, nitro, amino, sulfonamide, alkylamino, aralkyloxy, pyridyl, piperidino, carboxy, alkoxycarbonyl, acylamino, amino-

carbonyl, cyano or glucuronic acid residue; the cycloalkyl may be substituted or unsubstituted with a hydroxy, alkoxy or =O; the benzene-condensed cycloalkyl may be substituted or unsubstituted with a hydroxy or alkoxy; each of r, s, v and w independently represents 0 or 1; each of Y and Z independently represents a nitrogen atom, oxygen atom or sulfur atom; and each of p and q independently represents an integer of 1-4, provided that:


- (a) when R² is a hydrogen atom, then WR¹ is substituted at position-j of 2-oxoquinoline; and
 (b) 1,2-dihydro-6,7-dimethoxy-2-oxo-N-(phenylmethyl)-3-quinolinecarboxamide and N-(1,2-dihydro-6,7-dimethoxy-2-oxo-3-quinolyl) benzamide are excluded.

4. The 2-oxoquinoline compound of claim 3 or its pharmaceutically acceptable salt, wherein X' is -CONR^c-(Alk^a)_r-R.
 5. The 2-oxoquinoline compound of claim 3 or its pharmaceutically acceptable salt, wherein X' is -(CH₂)_p-OC(=Y)-NR^d-(Alk^b)_s-R or -(CH₂)_q-NR^e-C(=Z)-(NR^f)_w-(Alk^c)_v-R.
 6. The 2-oxoquinoline compound of any one of claims 3 to 5 or its pharmaceutically acceptable salt, wherein R is aryl, heteroaryl or



where aryl, heteroaryl, and each symbol in the formula are as defined above.

7. The 2-oxoquinoline compound of any one of claims 3 to 5 or its pharmaceutically acceptable salt, wherein R is



where each symbol is as defined above.

8. The 2-oxoquinoline compound of any one of claims 3 to 7 or its pharmaceutically acceptable salt, wherein W is -O- and R² is -OR⁶ (where R⁶ is a hydrogen atom or alkyl).
 9. The 2-oxoquinoline compound of any one of claims 3 to 8 or its pharmaceutically acceptable salt, wherein the substitution position of WR¹ is position-j of the benzene ring, and the substitution position of R² is position-i of the benzene ring.
 10. The 2-oxoquinoline compound of any one of claims 3, 4, and 6 to 9 or its pharmaceutically acceptable salt, wherein Alk^a is alkylene and r=1.
 11. The 2-oxoquinoline compound of any one of claims 3 to 10 or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminobenzyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-aminophenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-aminophenyl)amide hydrochloride, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-pyridylmethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-piperidinoethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-

3-carboxylic acid (2-morpholinoethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3-pyridylmethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridylmethyl)amide, 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide, 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide, 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide hydrochloride, 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxy-3-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-chloro-3-methoxyphenyl)ethyl]amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-methylbenzyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-fluorobenzyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide, 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide, 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid (2-phenylethyl)amide, 7,8-dimethoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide, 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-pyridine-4-ylethyl)amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-pyridylmethyl)amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-fluorobenzyl)amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-hydroxyphenyl)ethyl]amide, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (2-morpholinoethyl)amide, 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7,8-dipentyloxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 8-hydroxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (3,4-dihydroxybenzyl)amide, 1-O-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid (4-hydroxy-3-methoxybenzyl)amide, 1-O-2-hydroxy-5-[(7-methoxy-2-oxo-8-pentyloxy-1,2-dihydro-3-quinolyl)carbonylaminoethyl]phenyl]glucosidouronic acid, 1-O-2-hydroxy-4-[(7-methoxy-2-oxo-8-pentyloxy-1,2-dihydro-3-quinolyl)carbonylaminoethyl]phenyl]glucosidouronic acid, 5-[7-methoxy-3-[(3,4-methylenedioxybenzyl)carbonyl]-2-oxo-1,2-dihydro-8-quinolyl]pentanoic acid, 5-[7-methoxy-3-[(3-hydroxy-4-methoxybenzyl)carbonyl]-2-oxo-1,2-dihydro-8-quinolyl]pentanoic acid, 8-(5-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 8-(5-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (4-hydroxy-3-methoxybenzyl)amide, 8-(4-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7-methoxy-2-oxo-8-(4-oxopentyloxy)-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 8-(3-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7-methoxy-2-oxo-8-(3-oxopentyloxy)-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 8-(2-hydroxypentyloxy)-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid (3,4-methylenedioxybenzyl)amide, 7,8-dihydroxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid [2-(4-fluorophenyl)ethyl]amide, 8-butoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline, 8-ethoxy-3-hydroxymethyl-7-methoxy-2-oxo-1,2-dihydroquinoline, N-(4-fluorophenyl)carbamic acid (8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl ester, N-pyridine-4-ylcarbamic acid (8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl ester, 3-dimethylaminomethyl-8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline, 8-butoxy-3-aminomethyl-7-methoxy-2-oxo-1,2-dihydroquinoline, 8-ethoxy-7-methoxy-3-morpholinomethyl-2-oxo-1,2-dihydroquinoline, N-[(8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl]-N'-(4-fluorophenyl)urea and N-[(8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinolin-3-yl)methyl]-(4-hydroxyphenyl)acetamide.

12. A 2-oxoquinoline compound or its pharmaceutically acceptable salt, the 2-oxoquinoline compound being selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester, 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester, 1-methyl-7-methoxy-2-oxo-

8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester and 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid methyl ester.

13. A 2-oxoquinoline compound or its pharmaceutically acceptable salt, the 2-oxoquinoline compound being selected from the group consisting of 7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid, 8-butoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid, 8-ethoxy-7-methoxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acid, 7-methoxy-2-oxo-8-propoxy-1,2-dihydroquinoline-3-carboxylic acid, 7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid, 1-methyl-7-methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid and 1-methyl-7-methoxy-2-oxo-6-pentyloxy-1,2-dihydroquinoline-3-carboxylic acid.
14. 7-Methoxy-2-oxo-8-pentyloxy-1,2-dihydroquinoline-3-carboxamide or its pharmaceutically acceptable salt.
15. A pharmaceutical composition comprising, as an active ingredient, the 2-oxoquinoline compound of any one of claims 3 to 14 or its pharmaceutically acceptable salt.
16. A cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of any one of claims 3 to 14 or its pharmaceutically acceptable salt.
17. A peripheral cannabinoid receptor modulator comprising, as an active ingredient, the 2-oxoquinoline compound of any one of claims 3 to 14 or its pharmaceutically acceptable salt, the 2-oxoquinoline compound selectively acting on peripheral type cannabinoid receptors.
18. The 2-oxoquinoline compound of any one of claims 3 to 14 or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is an immunomodulator, therapeutic agent for autoimmune diseases, antiallergic agent, or antiinflammatory agent.
19. The 2-oxoquinoline compound of any one of claims 3 to 14 or its pharmaceutically acceptable salt, wherein the 2-oxoquinoline compound is an antiinflammatory agent.
20. An antiinflammatory agent comprising, as an active ingredient, the 2-oxoquinoline compound of claim 1 or 2 or its pharmaceutically acceptable salt.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/07398

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C07D215/48, 401/12, 405/12, C07H15/203, A61K31/47, 31/4709, 31/5355, 31/706, A61P37/02, 37/08, 29/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C07D215/48, 401/12, 405/12, C07H15/203 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN) REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EL-KERDAWY, M., et al., "Preparation of Alkoxyquinoline Derivatives and Their Evaluation as Potential Central Nervous System", Journal of Pharmaceutical Sciences, 73(11), 1652-3 (1984), Table I	3, 4, 6, 8, 10, 15
A		1, 2, 5, 7, 9, 11-14, 16-20
A	EP, 148623, A (Pfizer Cor.), 17 July, 1985 (17.07.85) & US, 4710507, A & ZA, 8409925, A & JP, 60-156687, A & HU, 36472, A & SU, 1407397, A	1-20
A	DE, 3134780, A (Hoechst A.G.), 10 March, 1983 (10.03.83) & EP, 73507, A & JP, 58-46068, A & ZA, 8206391, A & US, 4542139, A	1-20
PX	WO, 99/2499, A1 (JAPAN TOBACCO INC.), 21 January, 1999 (21.01.99) & AU, 9881279, A & JP, 11-80124, A	1-20
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
Date of the actual completion of the international search 07 April, 2000 (07.04.00)		Date of mailing of the international search report 18 April, 2000 (18.04.00)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)